

Constructing quantum vertex algebras

Haisheng Li¹

Department of Mathematical Sciences, Rutgers University, Camden, NJ 08102
and

Department of Mathematics, Harbin Normal University, Harbin, China

Abstract

This is a sequel to [Li6]. In this paper, we focus on the construction of quantum vertex algebras over $\mathbb{C}$, whose notion was formulated in [Li6] with Etingof and Kazhdan's notion of quantum vertex operator algebra (over $\mathbb{C}[[h]]$) as one of the main motivations. As one of the main steps in constructing quantum vertex algebras, we prove that every countable-dimensional nonlocal (namely noncommutative) vertex algebra over $\mathbb{C}$, which either is irreducible or has a basis of PBW type, is nondegenerate in the sense of Etingof and Kazhdan. Using this result, we establish the nondegeneracy of better known vertex operator algebras and some newly constructed nonlocal vertex algebras. We construct a family of quantum vertex algebras closely related to Zamolodchikov-Faddeev algebras.

1 Introduction

In a previous paper [Li6], to seek for appropriate vertex algebra-like objects for quantum affine algebras we studied fairly general formal vertex operators on an arbitrary vector space and we found that general formal vertex operators such as the generating functions of any quantum affine algebra in Drinfeld's realization (see [D]), acting on a highest weight module (see [FJ]), did give rise to vertex algebra-like structures in certain natural way. More specifically, given any vector space W (over $\mathbb{C}$), we studied what we called quasi compatible subsets of $\text{Hom}(W, W((x)))$ and we proved that every maximal quasi compatible subspace has a natural nonlocal (namely noncommutative) vertex algebra structure and that any quasi compatible subset generates a nonlocal vertex algebra. (Nonlocal vertex algebras were also called field algebras in [BK] and G_1 -vertex algebras in [Li3].) We found that for a quasi compatible set of formal vertex operators of a certain special type, the generated nonlocal vertex algebra very much resembles a quantum vertex operator algebra in the sense of Etingof and Kazhdan (see [EK]). This naturally led us to certain notions of weak quantum vertex algebra and quantum vertex algebra over $\mathbb{C}$.

Nonlocal vertex algebras are both vertex analogues and generalizations of noncommutative associative algebras and they are too general to work with. Naturally, one seeks for a good family of nonlocal vertex algebras which have more interesting structures. Then weak quantum vertex algebras are singled out for this consideration. Weak quantum vertex algebras are nonlocal vertex algebras (which are vertex analogues of noncommutative associative algebras), satisfying a certain generalized locality (where

¹Partially supported by an NSA grant

locality is a vertex analogue of classical commutativity). (Quantum vertex operator algebras in the sense of Etingof and Kazhdan are $\hbar$ -adic nonlocal vertex algebras satisfying a certain $\hbar$ -adic generalized locality.) Precisely, a weak quantum vertex algebra can be defined by using all the axioms in defining a vertex algebra V except the Jacobi identity axiom which is replaced by the following axiom: For any $u, v \in V$, there exists $\sum_{i=1}^r v^{(i)} \otimes u^{(i)} \otimes f_i(x) \in V \otimes V \otimes \mathbb{C}((x))$ such that

$$\begin{aligned} & x_0^{-1} \delta \left(\frac{x_1 - x_2}{x_0} \right) Y(u, x_1) Y(v, x_2) - x_0^{-1} \delta \left(\frac{x_2 - x_1}{-x_0} \right) \sum_{i=1}^r f_i(-x_0) Y(v^{(i)}, x_2) Y(u^{(i)}, x_1) \\ &= x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) Y(Y(u, x_0)v, x_2). \end{aligned} \quad (1.1)$$

Clearly, this notion generalizes the notions of vertex algebra, vertex superalgebra and vertex color algebra. A quantum vertex algebra is a weak quantum vertex algebra equipped with a “rational unitary quantum Yang-Baxter operator” $\mathcal{S}(x) : V \otimes V \rightarrow V \otimes V \otimes \mathbb{C}((x))$ such that for $u, v \in V$, (1.1) holds with

$$\sum_{i=1}^r v^{(i)} \otimes u^{(i)} \otimes f_i(x) = \mathcal{S}(x)(v \otimes u).$$

The notion of quantum vertex algebra clearly generalizes that of vertex algebra, vertex superalgebra and vertex color-algebra.

Just as ordinary vertex algebras can be constructed by using local sets of (formal) vertex operators (see [Li2]), it was proved in [Li6] that weak quantum vertex algebras can be constructed by using “ $\mathcal{S}$ -local” sets of (formal) vertex operators. Let W be any vector space. A subset U of $\text{Hom}(W, W((x)))$, alternatively denoted by $\mathcal{E}(W)$, is said to be $\mathcal{S}$ -local if for any $a(x), b(x) \in U$, there exists

$$\sum_{i=1}^r b^{(i)}(x) \otimes a^{(i)}(x) \otimes f_i(x) \in U \otimes U \otimes \mathbb{C}((x))$$

such that for some nonnegative integer k ,

$$(x_1 - x_2)^k a(x_1) b(x_2) = (x_1 - x_2)^k \sum_{i=1}^r f_i(x_2 - x_1) b^{(i)}(x_2) a^{(i)}(x_1), \quad (1.2)$$

where each $f_i(x_2 - x_1)$ is to be expanded in nonnegative powers of x_1 . For $a(x), b(x)$ from an $\mathcal{S}$ -local set U with the above information, we define $a(x)_n b(x) \in \mathcal{E}(W)$ for $n \in \mathbb{Z}$ as follows:

$$\begin{aligned} & a(x)_n b(x) \\ &= \text{Res}_{x_1} \left((x_1 - x_2)^n a(x_1) b(x_2) - \sum_{i=1}^r (-x_2 + x_1)^n f_i(x_2 - x_1) b^{(i)}(x_2) a^{(i)}(x_1) \right). \end{aligned} \quad (1.3)$$

Then it was proved that for any $\mathcal{S}$ -local subset U of $\mathcal{E}(W)$, there exists a unique closed $\mathcal{S}$ -local subspace $\langle U \rangle$ containing U and 1_W (the identity operator) and that $\langle U \rangle$ is a weak quantum vertex algebra with W naturally as a module. Furthermore, it is proved in [Li6] and this paper that if W and U are chosen to satisfy certain special properties, $\langle U \rangle$ and W are isomorphic $\langle U \rangle$ -modules. Consequently, W has a weak quantum vertex algebra structure transported from $\langle U \rangle$. This is analogous to a theorem of [FKRW] and [MP]. Next, to get a quantum vertex algebra from a weak quantum vertex algebra, one shall need to construct a unitary rational quantum Yang-Baxter operator with the desired property. For this we use an important idea of Etingof and Kazhdan.

In [EK], Etingof and Kazhdan introduced and studied a notion of quantum vertex operator algebra in the sense of formal deformation. By definition, the underlying space of a quantum vertex operator algebra V is a topologically free $\mathbb{C}[[h]]$ -module and the classical limit V/hV is an ordinary vertex algebra. Among the important ingredients of a quantum vertex operator algebra is a rational quantum Yang-Baxter operator $\mathcal{S}(x)$ (also depending on the formal parameter h). In that paper, they introduced a notion of nondegeneracy for a vertex algebra and proved that in constructing a quantum vertex operator algebra V (in the sense of [EK]), if V/hV (a vertex algebra over $\mathbb{C}$) is nondegenerate, the quantum Yang-Baxter relation follows from $\mathcal{S}$ -locality and associativity. An analogue which was obtained in [Li6] is that every nondegenerate weak quantum vertex algebra is a quantum vertex algebra.

Now we have seen that nondegeneracy of (ordinary) vertex algebras are very important for constructing quantum vertex operator algebras over $\mathbb{C}[[h]]$ in the sense of Etingof and Kazhdan while nondegeneracy of nonlocal vertex algebras are very important for constructing quantum vertex algebras over $\mathbb{C}$. The main purpose of this paper is to establish general results on nondegeneracy of nonlocal vertex algebras, which can be used to construct interesting examples of quantum vertex algebras.

A nonlocal vertex algebra V is *nondegenerate* if for each positive integer n the linear map $Z_n : V^{\otimes n} \otimes \mathbb{C}((x_1)) \cdots ((x_n)) \rightarrow V((x_1)) \cdots ((x_n))$, defined by

$$Z_n(u_1 \otimes \cdots \otimes u_n \otimes f) = fY(u_1, x_1) \cdots Y(u_n, x_n)\mathbf{1},$$

is injective. Let $\hat{\mathfrak{g}}$ be an affine Lie algebra and let ℓ be any complex number not equal to negative the dual Coxeter number of $\mathfrak{g}$. Associated to the pair $(\hat{\mathfrak{g}}, \ell)$ we have a vertex operator algebra $V_{\hat{\mathfrak{g}}}(\ell, 0)$ (cf. [FZ]), whose underlying space is the generalized Verma module or Weyl module of level ℓ for $\hat{\mathfrak{g}}$, induced from the 1-dimensional trivial $\mathfrak{g}$ -module $\mathbb{C}$. It was proved in [EK] that if $V_{\hat{\mathfrak{g}}}(\ell, 0)$ is an irreducible $\hat{\mathfrak{g}}$ -module, then the vertex operator algebra $V_{\hat{\mathfrak{g}}}(\ell, 0)$ is nondegenerate. Later, it was proved in [Li4] that for a general vertex algebra V , if V is of countable dimension over $\mathbb{C}$ and if the adjoint module V is irreducible, then V is nondegenerate. This in particular implies that every simple vertex operator algebra in the sense of [FLM] and [FHL] is nondegenerate.

In this paper, we first extend the result of [Li4], proving that if a nonlocal vertex algebra V is countable dimensional over $\mathbb{C}$ and if the adjoint module V is irreducible, then V is nondegenerate. Note that for certain rational numbers ℓ , even though the vertex operator algebras $V_{\hat{\mathfrak{g}}}(\ell, 0)$ are not simple, they play a very important role in

many studies. For the purpose of constructing quantum deformation of those nonsimple vertex operator algebras $V_{\mathfrak{g}}(\ell, 0)$ in the sense of [EK], we certainly hope that they are also nondegenerate. On the other hand, to apply the general construction theorem (Theorem 2.9) of quantum vertex algebras over $\mathbb{C}$, we often take the underlying space V to be a universal module for some algebra in a certain sense. (It seems that interesting nonlocal vertex algebras are most likely to be reducible, as most interesting noncommutative associative algebras are reducible under the left action and even not simple.) Motivated by this, we prove that every countable-dimensional nonlocal vertex algebra over $\mathbb{C}$, which is of PBW type in a certain sense, is nondegenerate. In particular, the vertex operator algebras $V_{\mathfrak{g}}(\ell, 0)$ associated to affine Lie algebras and those associated to the Virasoro Lie algebra are nondegenerate. This result enables us to construct quantum vertex algebras associated to certain subalgebras of affine Lie algebras. For example, for any finite-dimensional Lie algebra $\mathfrak{g}$, as d/dt is a derivation of the Lie algebra $t^{-1}\mathfrak{g}[t^{-1}]$, it can be extended uniquely to a derivation d of the universal enveloping algebra $U(t^{-1}\mathfrak{g}[t^{-1}])$. Then $U(t^{-1}\mathfrak{g}[t^{-1}])$ has a nonlocal vertex algebra structure with $Y(a, x)b = (e^{x^d}a)b$ for $a, b \in U(t^{-1}\mathfrak{g}[t^{-1}])$ ([B1], cf. [BK], [Li3]). We prove that this rather trivial nonlocal vertex algebra actually is a nondegenerate quantum vertex algebra with a nontrivial quantum Yang-Baxter operator. It is expected that from Yangians $Y(\mathfrak{g})$, which are closely related to $U(\mathfrak{g}[t])$, we have nondegenerate quantum vertex algebras. This will be addressed in a sequel.

Just as one needs associative or Lie algebras of a certain type such as affine Lie algebras and the Virasoro Lie algebra to construct interesting examples of vertex algebras, one needs algebras of a certain type to construct quantum vertex algebras. Indeed, there is a family of algebras, called Zamolodchikov-Faddeev algebras, which are very close to what we need. Zamolodchikov-Faddeev algebras are associative C^* -algebras, which have appeared in the study of quantum field theory (see [Z], [F], cf. [MR]). A Zamolodchikov-Faddeev algebra by definition is associated to an S -matrix, or a quantum Yang-Baxter operator (with two spectral parameters) on a finite-dimensional vector space U . Motivated by the construction of Zamolodchikov-Faddeev algebras, from any rational unitary quantum Yang-Baxter operator $\mathcal{S}(x) : H \otimes H \rightarrow H \otimes H \otimes \mathbb{C}[[x]]$ for a finite-dimensional vector space H with $\mathcal{S}(0) = 1$ we construct a vector space $V(H, \mathcal{S})$ and we prove that under a certain assumption if $V(H, \mathcal{S})$ is of PBW type in a certain sense, $V(H, \mathcal{S})$ has a unique nondegenerate quantum vertex algebra structure with certain properties. For $(H, \mathcal{S})$ of a certain special type, we show that indeed $V(H, \mathcal{S})$ is of PBW type.

Among our results we also refine the general construction theorem of [Li6] for weak quantum vertex algebras.

This paper is organized as follows: In Section 2 we review basic results on (weak) quantum vertex algebras and we prove certain new results. In Section 3, we establish certain basic results on nondegeneracy. In Section 4, we construct a family of quantum vertex algebras closely related to Zamolodchikov-Faddeev algebras.

2 Quantum vertex algebras and a general construction

In this section we first summarize some of the main results of [Li6] on weak quantum vertex algebras and quantum vertex algebras, and we then give a refinement of a construction theorem obtained in [Li6]. We also give an example to illustrate how to use the general construction theorem.

Throughout this paper we use the field of complex numbers $\mathbb{C}$ as the basic field. Following the tradition we denote by $\mathbb{C}(x_1, \dots, x_n)$ the field of rational functions. Set

$$\mathbb{C}_*(x_1, \dots, x_n) = \{f/g \mid f \in \mathbb{C}[[x_1, \dots, x_n]], 0 \neq g \in \mathbb{C}[x_1, \dots, x_n]\}, \quad (2.1)$$

a subring of the fraction field of $\mathbb{C}[[x_1, \dots, x_n]]$. So $\mathbb{C}_*(x_1, \dots, x_n)$ contains the field of rational functions $\mathbb{C}(x_1, \dots, x_n)$ as a subring. Denote by ι_{x_1, x_2} the natural embedding of $\mathbb{C}_*(x_1, x_2)$ into the field $\mathbb{C}((x_1))((x_2))$.

Throughout this paper, nonlocal vertex algebras are synonymous to G_1 -vertex algebras studied in [Li3] and they are also essentially field algebras studied in [BK] (cf. [K]).

Definition 2.1. A *nonlocal vertex algebra* is a vector space V equipped with a linear map

$$\begin{aligned} Y(\cdot, x) : V &\rightarrow \text{Hom}(V, V((x))) \subset (\text{End} V)[[x, x^{-1}]] \\ v &\mapsto Y(v, x) = \sum_{n \in \mathbb{Z}} v_n x^{-n-1} \quad (v_n \in \text{End} V) \end{aligned} \quad (2.2)$$

and equipped with a distinguished vector $\mathbf{1}$ such that for $v \in V$

$$Y(\mathbf{1}, x)v = v, \quad (2.3)$$

$$Y(v, x)\mathbf{1} \in V[[x]] \quad \text{and} \quad \lim_{x \rightarrow 0} Y(v, x)\mathbf{1} = v \quad (2.4)$$

and such that for $u, v, w \in V$, there exists a nonnegative integer l such that

$$(x_0 + x_2)^l Y(u, x_0 + x_2)Y(v, x_2)w = (x_0 + x_2)^l Y(Y(u, x_0)v, x_2)w. \quad (2.5)$$

Definition 2.2. A *V-module* is a vector space W equipped with a linear map

$$\begin{aligned} Y_W(\cdot, x) : V &\rightarrow \text{Hom}(W, W((x))) \subset (\text{End} W)[[x, x^{-1}]] \\ v &\mapsto Y_W(v, x) = \sum_{n \in \mathbb{Z}} v_n x^{-n-1} \quad (v_n \in \text{End} W) \end{aligned} \quad (2.6)$$

such that all the following axioms hold:

$$Y_W(\mathbf{1}, x) = 1_W \quad (\text{where } 1_W \text{ is the identity operator on } W), \quad (2.7)$$

and for any $u, v \in V$, $w \in W$, there exists $l \in \mathbb{N}$ such that

$$(x_0 + x_2)^l Y_W(u, x_0 + x_2)Y_W(v, x_2)w = (x_0 + x_2)^l Y_W(Y(u, x_0)v, x_2)w. \quad (2.8)$$

Definition 2.3. A *weak quantum vertex algebra* is a vector space V equipped with a vector $\mathbf{1}$ and a linear map $Y : V \rightarrow \text{Hom}(V, V((x)))$ such that for $v \in V$,

$$Y(\mathbf{1}, x)v = v, \quad (2.9)$$

$$Y(v, x)\mathbf{1} \in V[[x]] \quad \text{and} \quad \lim_{x \rightarrow 0} Y(v, x)\mathbf{1} = v \quad (2.10)$$

and such that for any $u, v \in V$, there exists $\sum_{i=1}^r v^{(i)} \otimes u^{(i)} \otimes f_i(x) \in V \otimes V \otimes \mathbb{C}((x))$ such that

$$\begin{aligned} & x_0^{-1} \delta \left(\frac{x_1 - x_2}{x_0} \right) Y(u, x_1)Y(v, x_2) - x_0^{-1} \delta \left(\frac{x_2 - x_1}{-x_0} \right) \sum_{i=1}^r f_i(-x_0)Y(v^{(i)}, x_2)Y(u^{(i)}, x_1) \\ &= x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) Y(Y(u, x_0)v, x_2). \end{aligned} \quad (2.11)$$

Remark 2.4. One can show that (2.11) is equivalent to the *$\mathcal{S}$ -locality*: For $u, v \in V$, there exists $\sum_{i=1}^r v^{(i)} \otimes u^{(i)} \otimes f_i(x) \in V \otimes V \otimes \mathbb{C}((x))$ such that

$$(x_1 - x_2)^k Y(u, x_1)Y(v, x_2) = (x_1 - x_2)^k \sum_{i=1}^r \iota_{x_2, x_1}(f_i(x_2 - x_1))Y(v^{(i)}, x_2)Y(u^{(i)}, x_1) \quad (2.12)$$

for some nonnegative integer k , and the *weak associativity*: For any $u, v, w \in V$, there exists a nonnegative integer l such that

$$(x_0 + x_2)^l Y(u, x_0 + x_2)Y(v, x_2)w = (x_0 + x_2)^l Y(Y(u, x_0)v, x_2)w. \quad (2.13)$$

Therefore a weak quantum vertex algebra is exactly a nonlocal vertex algebra V which satisfies $\mathcal{S}$ -locality.

Remark 2.5. Let V be a weak quantum vertex algebra and let (W, Y_W) be a module for V viewed as a nonlocal vertex algebra. From [Li6], whenever (2.11) holds, we have

$$\begin{aligned} & x_0^{-1} \delta \left(\frac{x_1 - x_2}{x_0} \right) Y_W(u, x_1)Y_W(v, x_2) \\ & \quad - x_0^{-1} \delta \left(\frac{x_2 - x_1}{-x_0} \right) \sum_{i=1}^r f_i(-x_0)Y_W(v^{(i)}, x_2)Y_W(u^{(i)}, x_1) \\ &= x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) Y_W(Y(u, x_0)v, x_2). \end{aligned} \quad (2.14)$$

In view of this, we simply define a V -module to be a module for V viewed as a nonlocal vertex algebra.

Let W be any vector space. Set

$$\mathcal{E}(W) = \text{Hom}(W, W((x))),$$

which in particular contains the identity operator 1_W on W . A subset U of $\mathcal{E}(W)$ is said to be $\mathcal{S}$ -local if for any $a(x), b(x) \in U$, there exist $f_i(x) \in \mathbb{C}((x))$, $b^{(i)}(x), a^{(i)}(x) \in U$ for $i = 1, \dots, r$, and a nonnegative integer k such that

$$(x_1 - x_2)^k a(x_1) b(x_2) = (x_1 - x_2)^k \sum_{i=1}^r \iota_{x_2, x_1}(f_i(x_2 - x_1)) b^{(i)}(x_2) a^{(i)}(x_1). \quad (2.15)$$

Notice that the above relation implies

$$(x_1 - x_2)^k a(x_1) b(x_2) \in \text{Hom}(W, W((x_1, x_2))). \quad (2.16)$$

Let U be any $\mathcal{S}$ -local subset of $\mathcal{E}(W)$. For $a(x), b(x) \in U$ and for $n \in \mathbb{Z}$ we define $a(x)_n b(x) \in \mathcal{E}(W)$ in terms of the generating function

$$Y_{\mathcal{E}}(a(x), x_0) b(x) = \sum_{n \in \mathbb{Z}} (a(x)_n b(x)) x_0^{-n-1} \quad (2.17)$$

by

$$Y_{\mathcal{E}}(a(x), x_0) b(x) = \text{Res}_{x_1} x^{-1} \delta \left(\frac{x_1 - x_0}{x} \right) x_0^{-k} ((x_1 - x)^k a(x_1) b(x)), \quad (2.18)$$

where k is any nonnegative integer such that $(x_1 - x_2)^k a(x_1) b(x_2) \in \text{Hom}(W, W((x_1, x_2)))$. Assuming the $\mathcal{S}$ -locality relation (2.15) we have

$$\begin{aligned} Y_{\mathcal{E}}(a(x), x_0) b(x) &= \text{Res}_{x_1} x_0^{-1} \delta \left(\frac{x_1 - x}{x_0} \right) a(x_1) b(x) \\ &\quad - \text{Res}_{x_1} x_0^{-1} \delta \left(\frac{x - x_1}{-x_0} \right) \sum_{i=1}^r \iota_{x, x_1}(f_i(x - x_1)) b^{(i)}(x) a^{(i)}(x_1), \end{aligned} \quad (2.19)$$

or equivalently, for $n \in \mathbb{Z}$,

$$\begin{aligned} &a(x)_n b(x) \\ &= \text{Res}_{x_1} \left((x_1 - x_2)^n a(x_1) b(x_2) - \sum_{i=1}^r (-x_2 + x_1)^n f_i(x_2 - x_1) b^{(i)}(x_2) a^{(i)}(x_1) \right) \end{aligned} \quad (2.20)$$

An $\mathcal{S}$ -local subspace U is said to be *closed* if $a(x)_n b(x) \in U$ for $a(x), b(x) \in U$, $n \in \mathbb{Z}$. The following is one of the results obtained in [Li6]:

Theorem 2.6. *Let W be any vector space and let U be any $\mathcal{S}$ -local subset of $\mathcal{E}(W)$. Then there exists a (unique) smallest closed $\mathcal{S}$ -local subspace $\langle U \rangle$, containing U and 1_W . Furthermore, $(\langle U \rangle, Y_{\mathcal{E}}, 1_W)$ is a weak quantum vertex algebra and (W, Y_W) carries the structure of a faithful V -module where $Y_W(\alpha(x), x_0) = \alpha(x_0)$ for $\alpha(x) \in V$.*

Let V be a nonlocal vertex algebra. A subset U of V is said to be $\mathcal{S}$ -local if $\{Y(u, x) \mid u \in U\}$ is an $\mathcal{S}$ -local subset of $\mathcal{E}(V)$. Note that a subset of an $\mathcal{S}$ -local subset is not necessarily $\mathcal{S}$ -local. Nevertheless we have:

Lemma 2.7. *Let V be a nonlocal vertex algebra and let U be an $\mathcal{S}$ -local subspace. Denote by $U^{(2)}$ the linear span of U and the vectors $a_m b$ for $a, b \in U$, $m \in \mathbb{Z}$. Then $U^{(2)}$ is $\mathcal{S}$ -local. Furthermore, the subalgebra $\langle U \rangle$ generated by U is a weak quantum vertex algebra.*

Proof. By assumption $\overline{U} = \{Y(v, x) \mid v \in V\}$ is an $\mathcal{S}$ -local subset of $\mathcal{E}(V)$. It follows from [Li6] (Lemmas 5.7 and 5.8) that $\overline{U}^{(2)}$ is $\mathcal{S}$ -local. For $u, v \in U$, by Proposition 6.6 of [Li6] (noting that $\mathcal{S}$ -locality implies compatibility), we have

$$Y(u_m v, x) = Y(u, x)_m Y(v, x).$$

It follows that $\overline{U^{(2)}} = \overline{U}^{(2)}$. Thus $\overline{U^{(2)}}$ is $\mathcal{S}$ -local. By definition $U^{(2)}$ is $\mathcal{S}$ -local.

For the second assertion, let K be a maximal $\mathcal{S}$ -local subspace of $\langle U \rangle$, containing U . Clearly, $K + \mathbb{C}\mathbf{1}$ is $\mathcal{S}$ -local. As K is maximal we have $\mathbf{1} \in K$. By the first assertion, $K^{(2)}$ is $\mathcal{S}$ -local. Again, as K is maximal we have $K^{(2)} = K$. Thus K is a subalgebra of $\langle U \rangle$, containing $\mathbf{1}$ and U . Therefore $\langle U \rangle = K$ and it is a weak quantum vertex algebra. $\square$

We shall need the following generalization of a result in [Li1] and [Li6]:

Proposition 2.8. *Let V be a weak quantum vertex algebra, let W be a V -module and let $e \in W$ be such that $Y(u, x)e \in W[[x]]$ for $u \in U$, where U is an $\mathcal{S}$ -local generating subspace of V . Then $Y(v, x)e \in W[[x]]$ for all $v \in V$. Furthermore, the linear map from V to W , sending v to $v_{-1}e$, is a V -module homomorphism.*

Proof. From [Li6] (cf. [Li1]), the second assertion is true if the first assertion is proved. Let F be the collection of $\mathcal{S}$ -local subspaces L containing U and satisfying the condition $Y(a, x)e \in W[[x]]$ for $a \in L$. By the assumption, $U \in F$. Let K be a maximal element of F . We are going to prove $K = V$. By assumption, $U \subset K$. Clearly $K + \mathbb{C}\mathbf{1} \in F$. It follows that $\mathbf{1} \in K$. As U generates V as a weak quantum vertex algebra, it suffices to prove that K is closed.

For any $a, b \in K$, there exist $a^{(i)}, b^{(i)} \in K$, $f_i(x) \in \mathbb{C}((x))$ for $i = 1, \dots, r$ such that

$$\begin{aligned} & x_2^{-1} \delta \left(\frac{x_1 - x_0}{x_2} \right) Y(Y(a, x_0)b, x_2)e - x_0^{-1} \delta \left(\frac{x_1 - x_2}{x_0} \right) Y(a, x_1)Y(b, x_2)e \\ &= -x_0^{-1} \delta \left(\frac{x_2 - x_1}{-x_0} \right) \sum_{i=1}^r \iota_{x_2, x_1}(f_i(x_2 - x_1)) Y(b^{(i)}, x_2) Y(a^{(i)}, x_1)e. \end{aligned} \quad (2.21)$$

As $Y(a^{(i)}, x_1)e \in W[[x_1]]$ for all the i 's, taking Res_{x_1} we get

$$Y(Y(a, x_0)b, x_2)e = \text{Res}_{x_1} x_0^{-1} \delta \left(\frac{x_1 - x_2}{x_0} \right) Y(a, x_1)Y(b, x_2)e = Y(a, x_0 + x_2)Y(b, x_2)e.$$

Since $Y(b, x_2)e \in W[[x_2]]$, the right-hand side lies in $W((x_0))[[x_2]]$. Consequently, we have

$$Y(Y(a, x_0)b, x_2)e \in W((x_2))((x_0)) \cap W((x_0))[[x_2]] \subset W[[x_0, x_2]][x_0^{-1}].$$

Thus $Y(a_mb, x)e \in W[[x]]$ for $m \in \mathbb{Z}$. On the other hand, by Lemma 2.7, $K^{(2)}$ is $\mathcal{S}$ -local. Therefore $K^{(2)} \in F$. Since K is maximal and $K \subset K^{(2)}$, we have $K^{(2)} = K$. This proves that K is closed and concludes the proof. $\square$

The following is our general construction theorem (cf. [Li6], [FKRW], [MP]):

Theorem 2.9. *Let V be a vector space, U a subset, $\mathbf{1}$ a vector of V , and Y_0 a map from U to $\mathcal{E}(V)$ ($= \text{Hom}(V, V((x)))$). Assume all the following conditions hold:*

$$Y_0(u, x)\mathbf{1} \in V[[x]] \quad \text{and} \quad \lim_{x \rightarrow 0} Y_0(u, x)\mathbf{1} = u \quad \text{for } u \in U, \quad (2.22)$$

$Y_0(U) = \{Y_0(u, x) \mid u \in U\} \subset \mathcal{E}(V)$ is $\mathcal{S}$ -local, and

$$V = \text{span}\{u_{n_1}^{(1)} \cdots u_{n_r}^{(r)}\mathbf{1} \mid r \geq 0, u^{(i)} \in U, n_i \in \mathbb{Z}\}, \quad (2.23)$$

where $Y_0(u, x) = \sum_{n \in \mathbb{Z}} u_n x^{-n-1}$ for $u \in U$. Furthermore, we assume that there exists a linear map f from V to $\langle Y_0(U) \rangle$ such that $f(\mathbf{1}) = 1_V$ and

$$f(u_n v) = u(x)_n f(v) \quad \text{for } u \in U, n \in \mathbb{Z}, v \in V. \quad (2.24)$$

Then Y_0 can be extended uniquely to a linear map Y from V to $\mathcal{E}(V)$ such that $(V, Y, \mathbf{1})$ carries the structure of a weak quantum vertex algebra.

Proof. We shall just prove the existence as the uniqueness is clear. By assumption, $Y_0(U)$ is an $\mathcal{S}$ -local subset of $\mathcal{E}(V)$. By Theorem 2.6, $Y_0(U)$ generates a weak quantum vertex algebra $\langle Y_0(U) \rangle$ inside $\mathcal{E}(V)$ with V as a module where $Y_V(\alpha(x), x_0) = \alpha(x_0)$ for $\alpha(x) \in \langle Y_0(U) \rangle$. With the assumption (2.22), by Proposition 2.8, the linear map g from $\langle Y_0(U) \rangle$ to V , sending $\alpha(x)$ to $\text{Res}_x x^{-1} \alpha(x)\mathbf{1}$, is a $\langle Y_0(U) \rangle$ -module homomorphism. Clearly, g sends 1_V to $\mathbf{1}$. Consequently, f and g are inverse each other. Now V has a weak quantum vertex algebra structure Y transported from $\langle Y_0(U) \rangle$ through the linear isomorphism f . For $u \in U$, as

$$f(u) = f(u_{-1}\mathbf{1}) = u(x)_{-1}1_V = u(x),$$

we have

$$Y(u, x_0) = gY_{\mathcal{E}}(u(x), x_0)f = Y_V(u(x), x_0) = u(x_0).$$

Indeed, Y extends the linear map Y_0 . This completes the proof. $\square$

Remark 2.10. In [Li6], a general construction theorem for nonlocal vertex algebras was given, which is an analogue of a construction theorem for vertex algebras of Frenkel-Kac-Ratiu-Wang [FKRW] and Meurman-Primc [MP]. In [Li6], $Y_0(U)$ was assumed to be quasi compatible (weaker than $\mathcal{S}$ -locality) and in addition it was assumed that there exists a linear operator $L(-1)$ on V such that

$$\begin{aligned} L(-1)\mathbf{1} &= 0, \\ [L(-1), Y_0(u, x)] &= \frac{d}{dx} Y_0(u, x) \quad \text{for } u \in U. \end{aligned}$$

The following result, obtained in [Li6], is an analogue of Theorem 5.7.6 of [LL]:

Theorem 2.11. *Let V be a weak quantum vertex algebra, let U be a generating subspace of V , let W be a vector space, and let Y_W^o be a linear map from U to $\mathcal{E}(W)$. Suppose that $Y_W^o(U)$ is $\mathcal{S}$ -local and that there exists a linear map f from V to $\langle Y_W^o(U) \rangle$ such that $f(\mathbf{1}) = 1_W$ and*

$$f(u_nv) = u(x)_n f(v) \quad \text{for } u \in U, n \in \mathbb{Z}, v \in V. \quad (2.25)$$

Then Y_W^o can be uniquely extended to a linear map Y_W from V to $\mathcal{E}(W)$ such that (W, Y_W) carries the structure of a V -module.

In practice we shall use the following result of [Li6] as a companion of Theorems 2.9 and 2.11:

Proposition 2.12. *Let W be a vector space, let V be a closed $\mathcal{S}$ -local subspace of $\mathcal{E}(W)$, and let*

$$u(x), v(x), u^{(1)}(x), v^{(1)}(x), \dots, u^{(r)}(x), v^{(r)}(x), c^0(x), \dots, c^s(x) \in V, \\ f_1(x), \dots, f_r(x) \in \mathbb{C}((x)).$$

Suppose that

$$(x_1 - x_2)^n u(x_1) v(x_2) - (-x_2 + x_1)^n \sum_{i=1}^r f_i(x_2 - x_1) v^{(i)}(x_2) u^{(i)}(x_1) \\ = \sum_{j=0}^s c^j(x_2) \frac{1}{j!} \left(\frac{\partial}{\partial x_2} \right)^j x_2^{-1} \delta \left(\frac{x_1}{x_2} \right) \quad (2.26)$$

for some integer n . Then

$$(x_1 - x_2)^n Y_{\mathcal{E}}(u(x), x_1) Y_{\mathcal{E}}(v(x), x_2) \\ - (-x_2 + x_1)^n \sum_{i=1}^r f_i(x_2 - x_1) Y_{\mathcal{E}}(v^{(i)}(x), x_2) Y_{\mathcal{E}}(u^{(i)}(x), x_1) \\ = \sum_{j=0}^s Y_{\mathcal{E}}(c^j(x), x_2) \frac{1}{j!} \left(\frac{\partial}{\partial x_2} \right)^j x_2^{-1} \delta \left(\frac{x_1}{x_2} \right). \quad (2.27)$$

Definition 2.13. A *quantum vertex algebra* is a weak quantum vertex algebra V equipped with a unitary rational quantum Yang-Baxter operator $\mathcal{S}$, which by definition is a linear map from $V \otimes V$ to $V \otimes V \otimes \mathbb{C}((x))$ satisfying the unitarity relation

$$\mathcal{S}_{21}(-x) \mathcal{S}(x) = 1 \quad (2.28)$$

and the following quantum Yang-Baxter equation

$$\mathcal{S}_{12}(x) \mathcal{S}_{13}(x+z) \mathcal{S}_{23}(z) = \mathcal{S}_{23}(z) \mathcal{S}_{13}(x+z) \mathcal{S}_{12}(x), \quad (2.29)$$

such that for $u, v \in V$, (2.11) holds with $\sum_{i=1}^r v^{(i)} \otimes u^{(i)} \otimes f_i(x) = \mathcal{S}(x)(v \otimes u)$.

The following result was proved in [Li6] (cf. [EK]):

Proposition 2.14. *Let V be a weak quantum vertex algebra. Assume that the linear map $\pi_3 : V \otimes V \otimes V \rightarrow \text{Hom}(V, V((x_1))((x_2))((x_3)))$, defined by*

$$\pi_3(u \otimes v \otimes w) = Y(u, x_1)Y(v, x_2)Y(w, x_3) \quad (2.30)$$

for $u, v, w \in V$, is injective. For $u, v \in V$, we define

$$\mathcal{S}(x)(v \otimes u) = \sum_{i=1}^r a_i \otimes b_i \otimes f_i(x) \in V \otimes V \otimes \mathbb{C}((x))$$

if

$$(x_1 - x_2)^k Y(u, x_1)Y(v, x_2) = (x_1 - x_2)^k \sum_{i=1}^r \iota_{x_2, x_1}(f_i(x_2 - x_1))Y(a_i, x_2)Y(b_i, x_1) \quad (2.31)$$

for some nonnegative integer k . Then $\mathcal{S}(x)$ is well defined and it is a rational unitary quantum Yang-Baxter operator on V and furthermore, $(V, \mathcal{S})$ is a quantum vertex algebra.

Closely related the linear map π_3 is the following notion of nondegeneracy due to Etingof and Kazhdan (see [EK]):

Definition 2.15. A nonlocal vertex algebra V is said to be *nondegenerate* if for each positive integer n the linear map $Z_n : V^{\otimes n} \otimes \mathbb{C}((x_1)) \cdots ((x_n)) \rightarrow V((x_1)) \cdots ((x_n))$, defined by

$$Z_n(u_1 \otimes \cdots \otimes u_n \otimes f) = fY(u_1, x_1) \cdots Y(u_n, x_n)1 \quad (2.32)$$

is injective.

Clearly, π_3 is injective whenever Z_3 is injective. Therefore we have:

Theorem 2.16. *Every nondegenerate weak quantum vertex algebra is a quantum vertex algebra where the quantum Yang-Baxter operator $\mathcal{S}(x)$ is uniquely determined.*

In the following we illustrate how to apply the general construction (Theorems 2.6 and 2.9 and Proposition 2.12) to construct weak quantum vertex algebras.

Definition 2.17. Let H be a vector space equipped with a bilinear form $\langle \cdot, \cdot \rangle$ and let $\mathcal{S}(x) : H \otimes H \rightarrow H \otimes H \otimes \mathbb{C}((x))$ be a linear map. An $(H, \mathcal{S})$ -module is a module W for the (free) tensor algebra $T(H \otimes \mathbb{C}[t, t^{-1}])$ such that for any $a \in H$, $w \in W$,

$$a(m)w = 0 \quad \text{for } m \text{ sufficiently large,} \quad (2.33)$$

where we use $a(m)$ for the operator corresponding to $a \otimes t^m$, and such that

$$a(x_1)b(x_2)w - \sum_{i=1}^r \iota_{x_2, x_1}(f_i(x_2 - x_1))b^{(i)}(x_2)a^{(i)}(x_1)w = x_2^{-1}\delta\left(\frac{x_1}{x_2}\right)\langle a, b \rangle w \quad (2.34)$$

for $a, b \in H$, $w \in W$, where $\mathcal{S}(x)(b \otimes a) = \sum_{i=1}^r b^{(i)} \otimes a^{(i)} \otimes f_i(x)$. (Notice that the condition (2.33) guarantees that for any $m, n \in \mathbb{Z}$, the coefficient of $x_1^m x_2^n$ in the left side is a finite sum in W .)

Let W be an $(H, \mathcal{S})$ -module. A vector $e \in W$ is called a *vacuum-like vector* if

$$a(m)e = 0 \quad \text{for all } a \in H, m \geq 0. \quad (2.35)$$

We call W a *vacuum $(H, \mathcal{S})$ -module* if W is cyclic on a vacuum-like vector.

Proposition 2.18. *Let H be a vector space equipped with a bilinear form $\langle \cdot, \cdot \rangle$ and let $\mathcal{S} : H \otimes H \rightarrow H \otimes H \otimes \mathbb{C}((x))$ be a linear map. Suppose that $V(H, \mathcal{S})$ is a universal vacuum $(H, \mathcal{S})$ -module with a vacuum vector $\mathbf{1}$ as its generator such that the linear map from H to $V(H, \mathcal{S})$, sending a to $a(-1)\mathbf{1}$ for $a \in H$, is injective. Then there exists a unique weak quantum vertex algebra structure on $V(H, \mathcal{S})$ with $\mathbf{1}$ as the vacuum vector such that*

$$Y(a(-1)\mathbf{1}, x) = a(x) \quad \text{for } a \in H. \quad (2.36)$$

Furthermore, on any $(H, \mathcal{S})$ -module W there exists a unique $V(H, \mathcal{S})$ -module structure Y_W such that

$$Y_W(a(-1)\mathbf{1}, x) = a(x) \quad \text{for } a \in H. \quad (2.37)$$

Proof. We are going to apply Theorem 2.9. Take $V = V(H, \mathcal{S})$ and $U = \{a(-1)\mathbf{1} \mid a \in H\}$. Define $Y_0(a(-1)\mathbf{1}, x) = a(x)$ for $a \in H$. The relation (2.34) implies that $\{Y_0(a(-1)\mathbf{1}, x) = a(x) \mid a \in H\}$ is $\mathcal{S}$ -local. Then all the assumptions except the last one (about the existence of a certain linear map f) hold.

For $a, b \in H$, with the relation (2.34) by Proposition 2.12 we have

$$\begin{aligned} & Y_{\mathcal{E}}(a(x), x_1)Y_{\mathcal{E}}(b(x), x_2) - \sum_{i=1}^r \iota_{x_2, x_1}(f_i(x_2 - x_1))Y_{\mathcal{E}}(b^{(i)}(x), x_2)Y_{\mathcal{E}}(a^{(i)}(x), x_1) \\ &= x_2^{-1}\delta\left(\frac{x_1}{x_2}\right)\langle a, b \rangle. \end{aligned}$$

Thus $\langle Y_0(U) \rangle$ is an $(H, \mathcal{S})$ -module with $a(x_0)$ acting as $Y_{\mathcal{E}}(a(x), x_0)$ for $a \in H$. Furthermore, $\langle Y_0(U) \rangle$ is a vacuum module with the vacuum vector 1_V as its generator. As $V(H, \mathcal{S})$ is a universal vacuum $(H, \mathcal{S})$ -module, there exists an $(H, \mathcal{S})$ -homomorphism f from $V(H, \mathcal{S})$ to $\langle Y_0(U) \rangle$, sending $\mathbf{1}$ to 1_V . Now it follows from Theorem 2.9. The last assertion follows from Theorem 2.11. $\square$

Remark 2.19. If the linear map π_3 defined in Proposition 2.14 for $V(H, \mathcal{S})$ is injective, it follows from Proposition 2.14 that $\mathcal{S}$ is necessarily a unitary rational quantum Yang-Baxter operator on H . In Section 4, we shall study this again after we establish certain results on nondegeneracy in Section 3.

Remark 2.20. The notion of $(H, \mathcal{S})$ -module was motivated by the following notion of Zamolodchikov-Faddeev algebra (see [Z], [F]). Let U be a finite-dimensional vector space over $\mathbb{C}$ and let $\mathcal{S}$ be a quantum Yang-Baxter operator on U , which by definition is a linear map

$$\mathcal{S}(x_1, x_2) : U \otimes U \rightarrow U \otimes U \otimes \mathbb{C}_*(x_1, x_2),$$

satisfying the quantum Yang-Baxter equation

$$\mathcal{S}_{12}(x_1, x_2)\mathcal{S}_{13}(x_1, x_3)\mathcal{S}_{23}(x_2, x_3) = \mathcal{S}_{23}(x_2, x_3)\mathcal{S}_{13}(x_1, x_3)\mathcal{S}_{12}(x_1, x_2). \quad (2.38)$$

The Zamolodchikov-Faddeev (ZF) algebra $\mathcal{A}(U, \mathcal{S})$ was defined to be an associative (C^*) -algebra with identity $\mathbf{1}$ with generators $\{a^{*\alpha}(\chi), a_\alpha(\chi) \mid \chi \in \mathbb{R}\}$ satisfying the following relations:

$$a_{\alpha_1}(\chi_1)a_{\alpha_2}(\chi_2) - \mathcal{S}_{\alpha_2\alpha_1}^{\beta_2\beta_1}(\chi_2, \chi_1)a_{\beta_2}(\chi_2)a_{\beta_1}(\chi_1) = 0, \quad (2.39)$$

$$a^{*\alpha_1}(\chi_1)a^{*\alpha_2}(\chi_2) - \mathcal{S}_{\beta_2\beta_1}^{\alpha_2\alpha_1}(\chi_2, \chi_1)a^{*\beta_2}(\chi_2)a^{*\beta_1}(\chi_1) = 0, \quad (2.40)$$

$$a_{\alpha_1}(\chi_1)a^{*\alpha_2}(\chi_2) - a^{*\beta_2}(\chi_2)\mathcal{S}_{\alpha_1\beta_2}^{\beta_1\alpha_2}(\chi_1, \chi_2)a_{\beta_1}(\chi_1) = 2\pi\delta_{\alpha_1\alpha_2}\delta(\chi_1 - \chi_2)\mathbf{1}. \quad (2.41)$$

We will not use this notion in this paper.

3 Nondegeneracy of nonlocal vertex algebras

Having known that every nondegenerate weak quantum vertex algebra is a quantum vertex algebra, in this section we determine when a nonlocal vertex algebra is nondegenerate. We prove that any irreducible nonlocal vertex algebra of countable dimension over $\mathbb{C}$ is nondegenerate, generalizing an earlier result of [Li4] from (ordinary) vertex algebras to nonlocal vertex algebras. We also prove that any nonlocal vertex algebra with a basis of PBW type is nondegenerate. Then we apply these results to establish the nondegeneracy of better known vertex (operator) algebras. We also construct some nondegenerate quantum vertex algebras associated to certain Lie algebras.

Recall that a nonlocal vertex algebra V is said to be nondegenerate if for each positive integer n the linear map $Z_n : V^{\otimes n} \otimes \mathbb{C}((x_1)) \cdots ((x_n)) \rightarrow V((x_1)) \cdots ((x_n))$, defined by

$$Z_n(u_1 \otimes \cdots \otimes u_n \otimes f) = fY(u_1, x_1) \cdots Y(u_n, x_n)\mathbf{1}$$

is injective.

Note that $\mathbb{C}((x_1)) \cdots ((x_n))$ is a field and $V^{\otimes n} \otimes \mathbb{C}((x_1)) \cdots ((x_n))$ and $V((x_1)) \cdots ((x_n))$ are vector spaces over $\mathbb{C}((x_1)) \cdots ((x_n))$. Clearly, the $\mathbb{C}$ -linear maps Z_n actually are $\mathbb{C}((x_1)) \cdots ((x_n))$ -linear maps.

First we have the following result:

Lemma 3.1. *Let U and V be nondegenerate nonlocal vertex algebras. Then the tensor product nonlocal vertex algebra $U \otimes V$ is nondegenerate.*

Proof. For any positive integer n , we have

$$\begin{aligned} & (U \otimes V)^{\otimes n} \otimes \mathbb{C}((x_1)) \cdots ((x_n)) \\ &= (U^{\otimes n} \otimes \mathbb{C}((x_1)) \cdots ((x_n))) \otimes_{\mathbb{C}((x_1)) \cdots ((x_n))} (V^{\otimes n} \otimes \mathbb{C}((x_1)) \cdots ((x_n))) \end{aligned}$$

and the image of $Z_n^{U \otimes V}$ (the Z_n map $U \otimes V$) is contained in

$$U((x_1)) \cdots ((x_n)) \otimes_{\mathbb{C}((x_1)) \cdots ((x_n))} V((x_1)) \cdots ((x_n)),$$

a subset of $(U \otimes V)((x_1)) \cdots ((x_n))$. Viewing Z_n^U, Z_n^V and $Z_n^{U \otimes V}$ as $\mathbb{C}((x_1)) \cdots ((x_n))$ -linear maps, we have

$$Z_n^{U \otimes V} = Z_n^U \otimes_{\mathbb{C}((x_1)) \cdots ((x_n))} Z_n^V.$$

As both Z_n^U and Z_n^V are injective, so is $Z_n^{U \otimes V}$. $\square$

Immediately from definition we have:

Lemma 3.2. *Let V be a nondegenerate nonlocal vertex algebra. Then any subalgebra of V is nondegenerate.*

On the other hand we have:

Lemma 3.3. *Let V be a nonlocal vertex algebra and let $\mathcal{D}$ be the linear operator defined by $\mathcal{D}(v) = v_{-2}\mathbf{1}$ for $v \in V$. If $\dim \ker \mathcal{D} \geq 2$, then V is degenerate. In particular, any associative algebra of dimension greater than 1 as a nonlocal vertex algebra is degenerate.*

Proof. We have $\mathbf{1} \in \ker \mathcal{D}$. If $\dim \ker \mathcal{D} \geq 2$, there exists $a \in \ker \mathcal{D}$ such that a and $\mathbf{1}$ are linearly independent. We have

$$0 \neq a \otimes \mathbf{1} \otimes \mathbf{1} - \mathbf{1} \otimes a \otimes \mathbf{1} \in V^{\otimes 2} \otimes \mathbb{C}((x_1))((x_2))$$

while

$$Z_2(a \otimes \mathbf{1} \otimes \mathbf{1} - \mathbf{1} \otimes a \otimes \mathbf{1}) = Y(a, x_1)\mathbf{1} - Y(a, x_2)\mathbf{1} = e^{x_1\mathcal{D}}a - e^{x_2\mathcal{D}}a = a - a = 0.$$

Thus Z_2 is not injective. $\square$

Example 3.4. Here we give some examples of degenerate vertex algebras. Let V be any vertex algebra and let (W, Y_W) be a V -module equipped with a linear operator d on W such that

$$[d, Y_W(v, x)] = \frac{d}{dx} Y_W(v, x) \quad \text{for } v \in V.$$

Extend the vertex operator map of V to the vector space $V \oplus W$ by

$$Y(u + w, x)(v + w') = Y(u, x)v + Y_W(u, x)w' + e^{xd}Y_W(v, -x)w$$

for $u, v \in V, w, w' \in W$. Then (cf. [Li1], [LL]) $V \oplus W$ is a vertex algebra with V a vertex subalgebra and with W an ideal. Since $Y(w, x)w' = 0$ for $w, w' \in W$, if $W \neq 0$, $V \oplus W$ is degenerate.

Let V be a nonlocal vertex algebra. A *quasi V -module* is a vector space W equipped with a linear map Y_W from V to $\mathcal{E}(W)$ such that $Y_W(\mathbf{1}, x) = 1_W$ and such that for any $u, v \in V$, $w \in W$,

$$p(x_0 + x_2, x_2)Y_W(u, x_0 + x_2)Y_W(v, x_2)w = p(x_0 + x_2, x_2)Y_W(Y(u, x_0)v, x_2)w \quad (3.1)$$

for some nonzero polynomial $p(x_1, x_2)$.

Definition 3.5. Let V be a nonlocal vertex algebra and let (W, Y_W) be a quasi V -module. For each positive integer n , we define a linear map

$$\pi_n^W : V^{\otimes n} \otimes \mathbb{C}((x_1)) \cdots ((x_n)) \rightarrow \text{Hom}(W, W((x_1)) \cdots ((x_n)))$$

by

$$\pi_n^W(u_1 \otimes \cdots \otimes u_n \otimes f) = f \cdot Y_W(u_1, x_1) \cdots Y_W(u_n, x_n). \quad (3.2)$$

Lemma 3.6. *Let V be a nonlocal vertex algebra and let n be a positive integer. If Z_n is injective, then π_n^V is injective. If V is a vertex algebra, the converse is also true.*

Proof. The first assertion is clear. Now, assume that V is a vertex algebra. Then for any $u, v \in V$, there exists a nonnegative integer k such that

$$(x_1 - x_2)^k Y(u, x_1)Y(v, x_2) = (x_1 - x_2)^k Y(v, x_2)Y(u, x_1).$$

In view of this, for any $a \in V$ and for any $A \in V^{\otimes n} \otimes \mathbb{C}((x_1)) \cdots ((x_n))$, there exists a nonnegative integer k such that

$$\left(\prod_{r=1}^n (x - x_r)^k \right) Y(a, x) \pi_n^V(A) = \left(\prod_{r=1}^n (x - x_r)^k \right) \pi_n^V(A) Y(a, x).$$

If $A \in \ker Z_n$, we have

$$\left(\prod_{r=1}^n (x - x_r)^k \right) \pi_n^V(A) Y(a, x) \mathbf{1} = \left(\prod_{r=1}^n (x - x_r)^k \right) Y(a, x) \pi_n^V(A) \mathbf{1} = 0.$$

Thus $\pi_n^V(A) Y(a, x) \mathbf{1} = 0$, which implies that $\pi_n^V(A)(a) = 0$. This proves that $A \in \ker \pi_n^V$. Therefore, if π_n^V is injective, so is Z_n . $\square$

Let $\hat{\mathfrak{g}} = \mathfrak{g} \otimes \mathbb{C}[t, t^{-1}] \oplus \mathbb{C}c$ be the affine Lie algebra associated to a (not necessarily finite-dimensional) Lie algebra $\mathfrak{g}$ equipped with a nondegenerate symmetric invariant bilinear form $\langle \cdot, \cdot \rangle$, where c is central and

$$[a \otimes t^m, b \otimes t^n] = [a, b] \otimes t^{m+n} + m \langle a, b \rangle \delta_{m+n, 0} c \quad (3.3)$$

for $a, b \in \mathfrak{g}$, $m, n \in \mathbb{Z}$. Associated to the affine Lie algebra $\hat{\mathfrak{g}}$ and any level $\ell \in \mathbb{C}$, we have a vertex algebra $V_{\hat{\mathfrak{g}}}(\ell, 0)$, which is the induced $\hat{\mathfrak{g}}$ -module from the 1-dimensional

$(\mathfrak{g}[t] \oplus \mathbb{C}c)$ -module $\mathbb{C}_\ell (= \mathbb{C})$ with $\mathfrak{g}[t]$ acting trivially and with c acting as the scalar ℓ . That is,

$$V_{\hat{\mathfrak{g}}}(\ell, 0) = U(\hat{\mathfrak{g}}) \otimes_{U(\mathfrak{g}[t] \oplus \mathbb{C}c)} \mathbb{C}_\ell \simeq U(t^{-1}\mathfrak{g}[t^{-1}]) = S(t^{-1}\mathfrak{g}[t^{-1}]). \quad (3.4)$$

This is an $\mathbb{N}$ -graded vertex algebra with $\mathbb{C}\mathbf{1}$ being the degree zero subspace and $\mathfrak{g}$, identified with the degree 1 subspace through the map $a \mapsto a(-1)\mathbf{1}$ for $a \in \mathfrak{g}$, generates $V_{\hat{\mathfrak{g}}}(\ell, 0)$ as a vertex algebra. If $\mathfrak{g}$ is finite-dimensional and simple, we often take $\langle \cdot, \cdot \rangle$ to be the normalized Killing form such that $\langle \alpha, \alpha \rangle = 2$ for long roots α . It is known (cf. [FZ], [DL]) that if ℓ is not equal to negative the dual Coxeter number of $\mathfrak{g}$, $V_{\hat{\mathfrak{g}}}(\ell, 0)$ is a vertex operator algebra (equipped with the Segal-Sugawara conformal vector). The following result was proved in [EK]:

Proposition 3.7. *If $V_{\hat{\mathfrak{g}}}(\ell, 0)$ is an irreducible $\hat{\mathfrak{g}}$ -module, then it is nondegenerate.*

It was proved in [Li4] that every irreducible vertex algebra of countable dimension over $\mathbb{C}$ is nondegenerate. Next we generalize this result from vertex algebras to nonlocal vertex algebras. First, we prove

Lemma 3.8. *Let V be a nonlocal vertex algebra and let (W, Y_W) be any quasi V -module. Let n be a positive integer and consider $V^{\otimes n} \otimes \mathbb{C}((x_1)) \cdots ((x_n))$ as a V -module with V acting on the first factor V . Then $\ker \pi_n^W$ is a V -submodule.*

Proof. Suppose that

$$X = \sum_i u_{1i} \otimes \cdots \otimes u_{ni} \otimes f_i(x_1, \dots, x_n) \in \ker \pi_n^W.$$

That is,

$$\sum_i f_i(x_1, \dots, x_n) Y_W(u_{1i}, x_1) \cdots Y_W(u_{ni}, x_n) = 0. \quad (3.5)$$

We must prove that for any $a \in V$ and any $w \in W$,

$$\sum_i f_i(x_1, \dots, x_n) Y_W(Y(a, x_0)u_{1i}, x_1) Y_W(u_{2i}, x_2) \cdots Y_W(u_{ni}, x_n) w = 0. \quad (3.6)$$

Let $m_2, \dots, m_n$ be arbitrarily fixed integers. Note that

$$\text{Res}_{x_2} \cdots \text{Res}_{x_n} x_2^{m_2} \cdots x_n^{m_n} f_i(x_1, \dots, x_n) Y_W(u_{2i}, x_2) \cdots Y_W(u_{ni}, x_n) w \in W \otimes \mathbb{C}((x_1))$$

for any i . Set

$$F_i = \text{Res}_{x_2} \cdots \text{Res}_{x_n} x_2^{m_2} \cdots x_n^{m_n} f_i(x_1, \dots, x_n) Y_W(u_{2i}, x_2) \cdots Y_W(u_{ni}, x_n) w.$$

For any $a \in V$, there exists $0 \neq p(x_1, x_2) \in \mathbb{C}[x_1, x_2]$ (depending on $m_2, \dots, m_n$) such that

$$p(x_0 + x_1, x_1) Y_W(a, x_0 + x_1) Y_W(u_{1i}, x_1) F_i = p(x_0 + x_1, x_1) Y_W(Y(a, x_0)u_{1i}x_1) F_i \quad (3.7)$$

for all the i . Then by applying $p(x_0 + x_1, x_1)Y_W(a, x_0 + x_1)\text{Res}_{x_2} \cdots \text{Res}_{x_n} x_2^{m_2} \cdots x_n^{m_n}$ to (3.5) we get

$$p(x_0 + x_1, x_1) \sum_i Y_W(Y(a, x_0)u_{1i}, x_1)F_i = 0. \quad (3.8)$$

By cancellation we have

$$\sum_i Y_W(Y(a, x_0)u_{1i}, x_1)F_i = 0. \quad (3.9)$$

That is,

$$\begin{aligned} & \text{Res}_{x_2} \cdots \text{Res}_{x_n} x_2^{m_2} \cdots x_n^{m_n} \sum_i f_i(x_1, \dots, x_n) Y_W(Y(a, x_0)u_{1i}, x_1) Y_W(u_{2i}, x_2) \cdots Y_W(u_{ni}, x_n) w \\ & = 0. \end{aligned} \quad (3.10)$$

Since $m_2, \dots, m_n$ are arbitrary, we have (3.6). This proves that $\ker \pi_n^W$ is a V -submodule. $\square$

Theorem 3.9. *Let V be an irreducible nonlocal vertex algebra of countable dimension over $\mathbb{C}$ and let (W, Y_W) be any quasi V -module. Then for each positive integer n , the linear map π_n^W is injective.*

Proof. Define π_0^W to be the natural embedding of $\mathbb{C}$ into $\text{Hom}(W, W)$. We shall prove that if π_n^W is not injective for some positive integer n , then π_{n-1}^W is not injective. Then the theorem follows immediately from induction.

Assume that π_n is not injective for some positive integer n . By Lemma 3.8, $\ker \pi_n^W$ is a V -submodule, where $V^{\otimes n} \otimes \mathbb{C}((x_1)) \cdots ((x_n))$ is considered as a V -module with V acting on the first factor. Let A be the associative algebra generated by operators v_m on $V^{\otimes n} \otimes \mathbb{C}((x_1)) \cdots ((x_n))$ for $v \in V$, $m \in \mathbb{Z}$. Since V is of countable-dimensional, we have $\text{End}_A V = \mathbb{C}$. Let $0 \neq X \in \ker \pi_n^W$. We have

$$X = u_1 \otimes e_1 + \cdots + u_r \otimes e_r,$$

where $u_1, \dots, u_r \in V$ are linearly independent and $0 \neq e_i \in V^{\otimes(n-1)} \otimes \mathbb{C}((x_1)) \cdots ((x_n))$. By Jacobson density theorem there exists $a \in A$ such that $au_1 = \mathbf{1}$ and $au_i = 0$ for $1 < i \leq r$. In view of Lemma 3.8 we have

$$aX = \mathbf{1} \otimes e_1 \in \ker \pi_n^W.$$

As $\mathbf{1} \otimes e_1 \in \ker \pi_n^W$, we have $\text{Res}_{x_n} x_n^r e_1 \in \ker \pi_{n-1}^W$ for $r \in \mathbb{Z}$. With $e_1 \neq 0$, $\text{Res}_{x_n} x_n^r e_1 \neq 0$ for some $r \in \mathbb{Z}$. Therefore, π_{n-1}^W is not injective. This completes the proof. $\square$

Corollary 3.10. *Every irreducible nonlocal vertex algebra of countable dimension over $\mathbb{C}$ is nondegenerate. In particular, simple vertex operator superalgebras over $\mathbb{C}$ (with finite-dimensional homogeneous subspaces) are nondegenerate.*

Remark 3.11. Let A be any unital (noncommutative) associative algebra and let ∂ be a derivation. Then ([B1], cf. [BK], [Li3]) A is a nonlocal vertex algebra with the identity 1 as the vacuum vector and with

$$Y(a, x)b = (e^{x\partial}a)b \quad \text{for } a, b \in A. \quad (3.11)$$

Let us denote this vertex algebra by (A, ∂) . Now take $A = \mathbb{C}[t]$ and $\partial = d/dt$. We have

$$Y(f(t), x)g(t) = (e^{x(d/dt)}f(t))g(t) = f(t+x)g(t) \quad \text{for } f(t), g(t) \in \mathbb{C}[t]. \quad (3.12)$$

It is easy to see that the vertex algebra $(\mathbb{C}[t], d/dt)$ is simple, but the adjoint module is reducible. Since

$$\begin{aligned} & Z_2(t \otimes 1 \otimes (x_1 - x_2)^{-1} - 1 \otimes t \otimes (x_1 - x_2)^{-1} - 1 \otimes 1 \otimes 1) \\ &= (x_1 - x_2)^{-1}(t + x_1) - (x_1 - x_2)^{-1}(t + x_2) - 1 \\ &= (x_1 - x_2)^{-1}(x_1 - x_2) - 1 \\ &= 0, \end{aligned}$$

Z_2 is not injective. This shows that the vertex algebra $(\mathbb{C}[t], d/dt)$ is degenerate. Thus the irreducibility assumption in Theorem 3.9 is necessary and it cannot be replaced by the simplicity. Furthermore, the degeneracy of vertex algebra $(\mathbb{C}[t], d/dt)$ implies that the vertex algebras $(\mathbb{C}[t, t^{-1}], d/dt)$ and $(\mathbb{C}((t)), d/dt)$ are degenerate as well. As $\mathbb{C}((t))$ is a field, $(\mathbb{C}((t)), d/dt)$ is irreducible. Thus the countability assumption in Theorem 3.9 is also necessary. Also, note that even though it is degenerate, we have $Y(f(t), x)g(t) = f(t+x)g(t) \neq 0$ for any $0 \neq f(t), g(t) \in \mathbb{C}((t))$.

Remark 3.12. For curiosity, one can show that $(\mathbb{C}[t^{-1}], d/dt)$ is nondegenerate.

Having proved that any irreducible nonlocal vertex algebra of countable dimension over $\mathbb{C}$ is nondegenerate, next we shall show that nonlocal vertex algebras with a basis of P-B-W type such as vertex algebras $V_{\hat{\mathfrak{g}}}(\ell, 0)$ are also nondegenerate. Our main tool will be increasing good filtrations which were studied in [Li5].

First we formulate the following straightforward result, which is classical in nature:

Lemma 3.13. *Let V be any nonlocal vertex algebra and let $E = \{E_n\}_{n \in \mathbb{Z}}$ be an increasing filtration of V satisfying the condition that $\mathbf{1} \in E_0$ and*

$$u_kv \in E_{m+n} \quad (3.13)$$

for $u \in E_m$, $v \in E_n$ and $k \in \mathbb{Z}$ with $m, n \in \mathbb{Z}$. Form the $\mathbb{Z}$ -graded vector space

$$\text{gr}_E(V) = \coprod_{n \in \mathbb{Z}} E_n / E_{n-1}. \quad (3.14)$$

Then $\text{gr}_E(V)$ is a nonlocal vertex algebra with $\mathbf{1} + E_{-1}$ as the vacuum vector and with

$$(u + E_{m-1})_k(v + E_{n-1}) = u_kv + E_{m+n-1} \quad (3.15)$$

for $u \in E_m$, $v \in E_n$ with $m, n, k \in \mathbb{Z}$.

We have:

Proposition 3.14. *Let V be a nonlocal vertex algebra. Assume that there exists a lower truncated increasing filtration $E = \{E_n\}$ satisfying the property (3.13) such that the associated nonlocal vertex algebra $\text{gr}_E(V)$ is nondegenerate. Then V is nondegenerate.*

Proof. First we construct a certain basis of V . By assumption, there is an integer r such that $E_n = 0$ for $n < r$. Let $\{v^\alpha \mid \alpha \in I_r\}$ be a basis of E_r , where I_r is an index set. We extend this basis of E_r to a basis $\{v^\alpha \mid \alpha \in I_r\} \cup \{v^\alpha \mid \alpha \in I_{r+1}\}$ of E_{r+1} , where I_{r+1} is another index set, and then extend this basis of E_{r+1} to a basis of E_{r+2} , and so on. In this way, we obtain a basis $\{v^\alpha \mid \alpha \in I\}$ of V , where $I = \cup_{m \geq r} I_m$ (a disjoint union). For $\alpha \in I_m$, we define $|\alpha| = m$. Then for $m \geq r$,

$$I_m = \{\alpha \in I \mid |\alpha| = m\}.$$

Let n be a positive integer. For $\alpha = (\alpha_1, \dots, \alpha_n) \in I^n$, set

$$|\alpha| = |\alpha_1| + \dots + |\alpha_n|.$$

From the construction of the basis $\{v^\alpha \mid \alpha \in I\}$, we have that $v^\alpha \in E_{|\alpha|}$ for $\alpha \in I$ and that for any $m \in \mathbb{Z}$, $\{v^\alpha + E_{m-1} \mid \alpha \in I_m\}$ is a basis of E_m/E_{m-1} .

Assume that Z_n for V is not injective. Then

$$\sum_{\alpha=(\alpha_1,\dots,\alpha_n) \in S} f_{(\alpha_1,\dots,\alpha_n)}(x_1, \dots, x_n) Y(v^{\alpha_1}, x_1) \cdots Y(v^{\alpha_n}, x_n) \mathbf{1} = 0 \quad (3.16)$$

for some finite subset S of I^n and $f_\alpha(x_1, \dots, x_n) \neq 0$ for $\alpha \in S$. Set $k = \max\{|\alpha| \mid \alpha \in S\}$. Then $S = S_r \cup S_{r+1} \cup \dots \cup S_k$.

From (3.13), for any $\alpha = (\alpha_1, \dots, \alpha_n) \in S_j$, we have

$$f_{(\alpha_1,\dots,\alpha_n)}(x_1, \dots, x_n) Y(v^{\alpha_1}, x_1) \cdots Y(v^{\alpha_n}, x_n) \mathbf{1} \in E_j((x_1)) \cdots ((x_n)).$$

It follows from (3.16) that

$$\begin{aligned} & \sum_{\alpha \in S_k} f_{(\alpha_1,\dots,\alpha_n)}(x_1, \dots, x_n) Y(v^{\alpha_1}, x_1) \cdots Y(v^{\alpha_n}, x_n) \mathbf{1} \\ &= - \sum_{m=r}^{k-1} \sum_{\alpha \in S_m} f_{(\alpha_1,\dots,\alpha_n)}(x_1, \dots, x_n) Y(v^{\alpha_1}, x_1) \cdots Y(v^{\alpha_n}, x_n) \mathbf{1} \\ &\in E_{k-1}((x_1)) \cdots ((x_n)). \end{aligned}$$

Thus

$$\sum_{\alpha \in S_k} f_{(\alpha_1,\dots,\alpha_n)}(x_1, \dots, x_n) Y(v^{\alpha_1} + E_{|\alpha_1|-1}, x_1) \cdots Y(v^{\alpha_n} + E_{|\alpha_n|-1}, x_n) \mathbf{1} = 0$$

in $(\text{gr}_E(V))^{\otimes n} \otimes \mathbb{C}((x_1)) \cdots ((x_n))$. From the construction of $\{v^\alpha \mid \alpha \in I\}$, we have

$$\sum_{\alpha \in S_k} (v^{\alpha_1} + E_{|\alpha_1|-1}) \otimes \cdots \otimes (v^{\alpha_n} + E_{|\alpha_n|-1}) \otimes f_{(\alpha_1, \dots, \alpha_n)}(x_1, \dots, x_n) \neq 0$$

in $(\text{gr}_E(V))^{\otimes n} \otimes \mathbb{C}((x_1)) \cdots ((x_n))$. This contradicts with our assumption that $\text{gr}_E(V)$ is nondegenerate. $\square$

Just as with classical associative algebras, for nonlocal vertex algebras we can obtain increasing filtrations from any generating subsets.

Proposition 3.15. *Let V be a nonlocal vertex algebra and let U be any generating subset of V . For any nonnegative integer n , define*

$$E_n^U = \text{span}\{u_{m_1}^{(1)} \cdots u_{m_r}^{(r)} \mathbf{1} \mid 0 \leq r \leq n, u^{(i)} \in U, m_i \in \mathbb{Z}\}. \quad (3.17)$$

Then $\{E_n^U\}_{n \in \mathbb{N}}$ is an increasing filtration of V satisfying the condition (3.13). Furthermore, if for $u, v \in U$, $m, n \in \mathbb{Z}$, and $k \geq 0$,

$$(u_m v_n - v_n u_m) E_k \subset E_{k+1}, \quad (3.18)$$

then the associated nonlocal vertex algebra $\text{gr}_E(V)$ is a (strictly) commutative vertex algebra.

Proof. As U generates V as a nonlocal vertex algebra, $\{E_n^U\}_{n \in \mathbb{N}}$ is an increasing filtration of V . To establish the property (3.13), we are going to use induction to prove that for any $m \geq 0$,

$$a_k E_n \subset E_{m+n}$$

for all $a \in E_m$, $k \in \mathbb{Z}$. Since $E_0 = \mathbb{C}\mathbf{1}$, it is true for $m = 0$. Now assume $a \in E_{m+1}$. Notice that for any $m \geq 0$,

$$E_{m+1} = \text{span}\{u_r v \mid u \in U, r \in \mathbb{Z}, v \in E_m\}.$$

In view of this, it suffices to consider $a = u_r v$ for $u \in U$, $r \in \mathbb{Z}$, $v \in E_m$. For $w \in E_n$, there exists a nonnegative integer l such that

$$(x_0 + x_2)^l Y(Y(u, x_0)v, x_2)w = (x_0 + x_2)^l Y(u, x_0 + x_2)Y(v, x_2)w.$$

With $v \in E_m$, $w \in E_n$, using inductive hypothesis we have

$$Y(u, x_0 + x_2)Y(v, x_2)w \in Y(u, x_0 + x_2)E_{n+m}((x_2)) \subset E_{n+m+1}((x_0))((x_2)).$$

Consequently we have

$$(x_0 + x_2)^l Y(Y(u, x_0)v, x_2)w \in E_{n+m+1}((x_0))((x_2)).$$

Since $Y(Y(u, x_0)v, x_2)w \in V((x_2))((x_0))$, we have

$$Y(Y(u, x_0)v, x_2)w \in E_{n+m+1}((x_2))((x_0)).$$

In particular, we have $a_k E_n \subset E_{n+m+1}$.

Notice that $U \subset E_1$. As U generates V as a nonlocal vertex algebra, U/E_0 generates $\text{gr}_E(V)$ as a nonlocal vertex algebra. Then the last assertion is clear. $\square$

We next recall certain results on constructing increasing filtrations for vertex algebras from [Li5]:

Lemma 3.16. *Let V be an (ordinary) vertex algebra. Let $E = \{E_n\}$ be an increasing good filtration of V in the sense that $\mathbf{1} \in E_0$ and that for $u \in E_m$, $v \in E_n$*

$$u_k v \in E_{m+n} \quad \text{for any } k \in \mathbb{Z},$$

and $u_k v \in E_{m+n-1}$ for $k \geq 0$. Then $\text{gr}_E(V)$ is a strictly commutative vertex algebra, which amounts to a unital commutative associative differential algebra.

Theorem 3.17. *Let V be an (ordinary) vertex algebra and let U be a subspace of V equipped with a vector space decomposition $U = \coprod_{n \geq 1} U_n$. For $n \geq 0$, denote by E_n^U the subspace of V linearly spanned by the vectors*

$$u_{-k_1}^{(1)} \cdots u_{-k_r}^{(r)} \mathbf{1}$$

for $r \geq 0$, $u^{(i)} \in U$, $k_i \geq 1$ with $\text{wt } u^{(1)} + \cdots + \text{wt } u^{(r)} \leq n$. Assume that U generates V as a vertex algebra and that for $u \in U_m$, $v \in U_n$ with $m, n \geq 1$, $u_i v \in E_{m+n-1}$ for all $i \geq 0$. Then $\{E_n^U\}$ is an increasing good filtration of V and $\text{gr}_{E^U}(V)$ is a (strictly) commutative vertex algebra. Furthermore, any good filtration $\{E_n\}$ of V with $E_n = 0$ for $n < 0$ and with $E_0 = \mathbb{C}\mathbf{1}$ can be obtained this way.

Next, we establish nondegeneracy for better known vertex operator algebras. Let U be any vector space over $\mathbb{C}$. Set

$$FCD(U) = S(\mathbb{C}[\partial] \otimes U), \tag{3.19}$$

the symmetric algebra over the vector space $\mathbb{C}[\partial] \otimes U$, where ∂ is a formal variable. Then $FCD(U)$ is a commutative associative algebra equipped with a derivation ∂ . We call $FCD(U)$ the *free commutative differential algebra over U* .

Proposition 3.18. *Let U be any vector space of countable dimension over $\mathbb{C}$. The free commutative differential algebra $FCD(U)$ on U , viewed as a vertex algebra, is nondegenerate.*

Proof. Pick up a basis $\{u_\alpha \mid \alpha \in I\}$ of U . Let U' be the subspace of U^* , linearly spanned by the linear functionals u_α^* dual to u_α for $\alpha \in I$. Set $\mathfrak{h} = U \oplus U'$, a vector space of countable dimension over $\mathbb{C}$. Equip $\mathfrak{h}$ with the standard (nondegenerate) symmetric bilinear form $\langle \cdot, \cdot \rangle$

$$\langle a + a', b + b' \rangle = \langle a, b' \rangle + \langle a', b \rangle$$

for $a, b \in U$, $a', b' \in U'$. As $V_{\mathfrak{h}}(1, 0)$ is an irreducible vertex algebra of countable dimension over $\mathbb{C}$, by Corollary 3.10, $V_{\mathfrak{h}}(1, 0)$ is nondegenerate. Consider the vertex

subalgebra $\langle U \rangle$ of $V_{\mathfrak{h}}(1, 0)$, generated by the subspace U . In view of Lemma 3.2, $\langle U \rangle$ is nondegenerate. For $a, b \in U$ and for $m, n \in \mathbb{Z}$, we have

$$[a(m), b(n)] (= [a, b](m+n) + m\langle a, b \rangle \delta_{m+n, 0}) = 0.$$

As U generates $\langle U \rangle$ as a vertex algebra, $\langle U \rangle$ is a (strictly) commutative vertex algebra. Equivalently, $\langle U \rangle$ is a commutative differential algebra. In view of the P-B-W theorem, we have $\langle U \rangle = S(t^{-1}U[t^{-1}])$ as a vector space. Consequently, we have that $FCD(U) = \langle U \rangle$ as a commutative differential algebra and then as a vertex algebra. Therefore, $FCD(U)$ is nondegenerate. $\square$

Combining Propositions 3.14 and 3.18 we immediately have:

Theorem 3.19. *Let V be a nonlocal vertex algebra of countable dimension over $\mathbb{C}$. Assume that there exists an increasing filtration $E = \{E_n\}_{n \geq 0}$ of V , satisfying the condition (3.13), such that the associated nonlocal vertex algebra $\text{gr}_E(V)$ is isomorphic to the free commutative differential algebra $FCD(U)$ over a vector space U . Then V is nondegenerate.*

In view of Theorem 2.16 we immediately have:

Corollary 3.20. *Let V be a weak quantum vertex algebra of countable dimension over $\mathbb{C}$. Assume that there exists an increasing filtration $E = \{E_n\}_{n \geq 0}$ of V , satisfying the condition (3.13), such that the associated nonlocal vertex algebra $\text{gr}_E(V)$ is isomorphic to the free commutative differential algebra $FCD(U)$ over a vector space U . Then V is a nondegenerate quantum vertex algebra.*

Now we are ready to establish the nondegeneracy of vertex algebra $V_{\mathfrak{g}}(\ell, 0)$ in the general case (cf. [EK]).

Proposition 3.21. *Let $\mathfrak{g}$ be any Lie algebra of countable dimension over $\mathbb{C}$ equipped with a nondegenerate symmetric invariant bilinear form. Then for any complex number ℓ , the vertex algebra $V_{\mathfrak{g}}(\ell, 0)$ is nondegenerate.*

Proof. Recall that $V_{\mathfrak{g}}(\ell, 0)$ as a vertex algebra is generated by $\mathfrak{g}$. Taking $U = U_1 = \mathfrak{g}$ in Theorem 3.17 we obtain a good increasing filtration $E^{\mathfrak{g}} = \{E_n^{\mathfrak{g}}\}_{n \geq 0}$ of $V_{\mathfrak{g}}(\ell, 0)$. For any positive integer n , $E_n^{\mathfrak{g}}$ is the linear span of the vectors

$$a^{(1)}(-m_1) \cdots a^{(r)}(-m_r) \mathbf{1}$$

for $0 \leq r \leq n$, $a^{(1)}, \dots, a^{(r)} \in \mathfrak{g}$, $m_1, \dots, m_r \geq 1$. It follows from the P-B-W theorem that $V_{\mathfrak{g}}(\ell, 0) = U(t^{-1}\mathfrak{g}[t^{-1}])$ as a vector space. The filtration E coincides with the standard filtration on $U(t^{-1}\mathfrak{g}[t^{-1}])$. Then $\text{gr}_E(V_{\mathfrak{g}}(\ell, 0)) = S(t^{-1}\mathfrak{g}[t^{-1}])$. Thus $\text{gr}_E(V_{\mathfrak{g}}(\ell, 0))$, a (strictly) commutative vertex algebra, is isomorphic to the free commutative differential algebra over $\mathfrak{g}$. By Theorem 3.19, $V_{\mathfrak{g}}(\ell, 0)$ is nondegenerate. $\square$

Consider the Virasoro Lie algebra $Vir = \sum_{n \in \mathbb{Z}} \mathbb{C}L(n) \oplus \mathbb{C}c$. Recall (cf. [FZ]) that for any complex number ℓ , we have a vertex operator algebra $V_{Vir}(\ell, 0)$, associated to the Virasoro Lie algebra Vir of central charge ℓ . By definition,

$$V_{Vir}(\ell, 0) = U(Vir) \otimes_{U(Vir_{(\geq 0)})} \mathbb{C}_\ell, \quad (3.20)$$

where $\mathbb{C}_\ell$ is the 1-dimensional module for the Lie subalgebra

$$Vir_{(\geq 0)} = \sum_{n \geq 0} \mathbb{C}L(n-1) \oplus \mathbb{C}c$$

with c acting as scalar ℓ and with $\sum_{n \geq 0} \mathbb{C}L(n-1)$ acting as zero.

Proposition 3.22. *For any complex number ℓ , the vertex operator algebra $V_{Vir}(\ell, 0)$ is nondegenerate.*

Let $\mathcal{L}$ be a Lie algebra equipped with a derivation d . Then d naturally extends to a derivation of $U(\mathcal{L})$, the universal enveloping algebra of $\mathcal{L}$. From Remark 3.11, $U(\mathcal{L})$ is naturally a nonlocal vertex algebra.

Proposition 3.23. *Let $\mathcal{L}$ be a Lie algebra equipped with a derivation d such that $\mathcal{L}$ is a free $\mathbb{C}[d]$ -module with a countable base. Then the nonlocal vertex algebra $U(\mathcal{L})$ is nondegenerate.*

Proof. Recall that for $a, b \in U(\mathcal{L})$ we have $Y(a, x)b = (e^{xd}a)b$. That is, $a_nb = 0$ for $n \geq 0$ and $a_nb = \frac{1}{(-n-1)!}(d^{-n-1}a)b$ for $n < 0$. As $\mathcal{L}$ generates $U(\mathcal{L})$ as an algebra, $\mathcal{L}$ generates $U(\mathcal{L})$ as a nonlocal vertex algebra. Also, for the increasing filtration E defined in Proposition 3.15 we have

$$E_n = \text{span}\{a^{(1)} \cdots a^{(r)} \mid 0 \leq r \leq n, a^{(i)} \in \mathcal{L}\}$$

for $n \geq 0$. Thus, the filtration E is exactly the standard filtration for the universal enveloping algebra $U(\mathcal{L})$. It was well known that $\text{gr}_E(U(\mathcal{L})) = S(\mathcal{L})$ (the symmetric algebra). By Theorem 3.19 the nonlocal vertex algebra $U(\mathcal{L})$ is nondegenerate. $\square$

We next consider current Lie algebras. Let $\mathfrak{g}$ be any Lie algebra. With $t^{-1}\mathbb{C}[t^{-1}]$ being a commutative associative algebra, $\mathfrak{g} \otimes t^{-1}\mathbb{C}[t^{-1}]$ is naturally a Lie algebra. Furthermore, $L(-1) = 1 \otimes d/dt$ is a derivation. It is clear that $\mathfrak{g} \otimes t^{-1}\mathbb{C}[t^{-1}]$ is a free $\mathbb{C}[L(-1)]$ -module. In view of Proposition 3.23 we immediately have:

Corollary 3.24. *Let $\mathfrak{g}$ be any Lie algebra of countable dimension over $\mathbb{C}$. Then the nonlocal vertex algebra $U(\mathfrak{g} \otimes t^{-1}\mathbb{C}[t^{-1}])$ associated to the noncommutative differential algebra structure is nondegenerate.*

Furthermore we have:

Proposition 3.25. *Let $\mathfrak{g}$ be any Lie algebra of countable dimension over $\mathbb{C}$. Then the nonlocal vertex algebra $U(\mathfrak{g} \otimes t^{-1}\mathbb{C}[t^{-1}])$ associated to the noncommutative differential algebra structure is a nondegenerate quantum vertex algebra.*

Proof. For $a \in \mathfrak{g}$, set

$$a(x)^+ = \sum_{n \geq 1} (a \otimes t^{-n}) x^{n-1} \in (\mathfrak{g} \otimes t^{-1} \mathbb{C}[t^{-1}])[[x]]. \quad (3.21)$$

We have

$$Y(a \otimes t^{-1}, x) = e^{xL(-1)}(a \otimes t^{-1}) = a \otimes (t+x)^{-1} = \sum_{n \geq 0} (a \otimes t^{-n-1}) x^n = a(x)^+.$$

It is clear that $\mathfrak{g} \otimes t^{-1}$ generates $U(\mathfrak{g} \otimes t^{-1} \mathbb{C}[t^{-1}])$ as a nonlocal vertex algebra. For $a, b \in \mathfrak{g}$, by a straightforward calculation we have

$$[a(x_1)^+, b(x_2)^+] = (x_2 - x_1)^{-1} ([a, b](x_2)^+ - [a, b](x_1)^+). \quad (3.22)$$

From this, we have

$$(x_1 - x_2)a(x_1)^+b(x_2)^+ = (x_1 - x_2)b(x_2)^+a(x_1)^+ + [a, b](x_2)^+ \cdot 1 - 1 \cdot [a, b](x_1)^+.$$

This shows that $\{Y(a \otimes t^{-1}, x) = a(x)^+ \mid a \in \mathfrak{g}\}$ is $\mathcal{S}$ -local and that $\mathfrak{g} \otimes t^{-1}$ is an $\mathcal{S}$ -local generating subspace of $U(\mathfrak{g} \otimes t^{-1} \mathbb{C}[t^{-1}])$. By Lemma 2.7, $U(\mathfrak{g} \otimes t^{-1} \mathbb{C}[t^{-1}])$ is a weak quantum vertex algebra. As it has been proved in Corollary 3.24 that $U(\mathfrak{g} \otimes t^{-1} \mathbb{C}[t^{-1}])$ is nondegenerate, the rest follows from Theorem 2.16. $\square$

Remark 3.26. Let $\mathfrak{g}$ be a Lie algebra of countable dimension over $\mathbb{C}$. Set $U = \mathfrak{g} \oplus \mathbb{C}$. It follows from Proposition 3.25, we have a unitary rational quantum Yang-Baxter operator $\mathcal{S}$ on U such that

$$\mathcal{S}(x)(1 \otimes 1) = 1 \otimes 1, \quad (3.23)$$

$$\mathcal{S}(x)(1 \otimes a) = 1 \otimes a, \quad (3.24)$$

$$\mathcal{S}(x)(a \otimes 1) = a \otimes 1, \quad (3.25)$$

$$\mathcal{S}(x)(a \otimes b) = a \otimes b + x^{-1}(\mathbf{1} \otimes [a, b] - [a, b] \otimes \mathbf{1}) \quad (3.26)$$

for $a, b \in \mathfrak{g}$.

Remark 3.27. Consider the Lie algebra $\mathfrak{g}[t]$. Let d be any derivation of the polynomial algebra $\mathbb{C}[t]$. Then d is naturally a derivation of the Lie algebra $\mathfrak{g}[t]$, hence a derivation of $U(\mathfrak{g}[t])$. Now we have a nonlocal vertex algebra $U(\mathfrak{g}[t])$. But, since $\mathcal{D} = d$ and $\mathfrak{g} \oplus \mathbb{C} \subset \ker d$, by Lemma 3.3 $U(\mathfrak{g}[t])$ is degenerate.

Remark 3.28. Note that Yangian $Y(\mathfrak{g})$ is closely related to the Lie algebra $\mathfrak{g}[t]$. In fact, the associated graded algebra of $Y(\mathfrak{g})$ with respect to a certain increasing filtration is isomorphic to $U(\mathfrak{g}[t])$. We shall study Yangian $Y(\mathfrak{g})$ in terms of quantum vertex algebras in a separate publication.

Next we construct some examples of nondegenerate quantum vertex algebras from certain Lie algebras. Consider a semidirect product Lie algebra $K \oplus \mathfrak{g}$, where K is an ideal and $\mathfrak{g}$ is a Lie subalgebra. Suppose that $\langle \cdot, \cdot \rangle$ is a symmetric invariant bilinear form on K . Extend $\langle \cdot, \cdot \rangle$ to a bilinear form on $K \oplus \mathfrak{g}$ by $\langle u + a, v + b \rangle = \langle u, v \rangle$ for $u, v \in K$, $a, b \in \mathfrak{g}$. We have an affine Lie algebra associated to $(K \oplus \mathfrak{g}, \langle \cdot, \cdot \rangle)$. Consider the following Lie subalgebra

$$\mathcal{L} = K \otimes \mathbb{C}[t, t^{-1}] \oplus \mathbb{C}c \oplus (\mathfrak{g} \otimes t^{-1}\mathbb{C}[t^{-1}]), \quad (3.27)$$

which is the semidirect product Lie algebra of $\hat{K}$ with $\mathfrak{g} \otimes t^{-1}\mathbb{C}[t^{-1}]$. Let ℓ be any complex number. Let $\mathbb{C}_\ell$ be the 1-dimensional module for $K[t] \oplus \mathbb{C}c$ with $K[t]$ acting as zero and with c acting as scalar ℓ . Form the induced $\mathcal{L}$ -module

$$V_{\mathcal{L}}(\ell, 0) = U(\mathcal{L}) \otimes_{U(K[t] \oplus \mathbb{C}c)} \mathbb{C}_\ell. \quad (3.28)$$

In view of the P-B-W theorem, we have

$$V_{\mathcal{L}}(\ell, 0) = U((K \oplus \mathfrak{g}) \otimes t^{-1}\mathbb{C}[t^{-1}]), \quad (3.29)$$

as a vector space. Embed $K \oplus \mathfrak{g}$ into $V_{\mathcal{L}}(\ell, 0)$ through the map $u + a \mapsto u(-1)\mathbf{1} + a(-1)\mathbf{1}$.

Proposition 3.29. *There exists a unique weak quantum vertex algebra structure on $V_{\mathcal{L}}(\ell, 0)$ such that*

$$Y(u + a, x) = u(x) + a(x)^+ \quad \text{for } u \in K, a \in \mathfrak{g}. \quad (3.30)$$

Furthermore, $V_{\mathcal{L}}(\ell, 0)$ is a nondegenerate quantum vertex algebra.

Proof. We shall apply Theorem 2.9. Set

$$U = \{u(x), a(x)^+ \mid u \in K, a \in \mathfrak{g}\}.$$

We now prove that U is $\mathcal{S}$ -local. For $u, v \in K$, we have

$$[u(x_1), v(x_2)] = [u, v](x_2)x_1^{-1}\delta\left(\frac{x_2}{x_1}\right) + \ell\langle u, v \rangle \frac{\partial}{\partial x_2} x_1^{-1}\delta\left(\frac{x_2}{x_1}\right).$$

For $u \in K$, $a \in \mathfrak{g}$, we have

$$[a(x_1)^+, u(x_2)] = [a, u](x_2)(x_2 - x_1)^{-1}.$$

For $a, b \in \mathfrak{g}$, we have

$$[a(x_1)^+, b(x_2)^+] = (x_2 - x_1)^{-1}([a, b](x_2)^+ - [a, b](x_1)^+).$$

It follows that U is $\mathcal{S}$ -local. By Theorem 2.6, U generates a weak quantum vertex algebra $\langle U \rangle$ inside $\mathcal{E}(V)$. It follows from Proposition 2.12 that $\langle U \rangle$ is naturally an $\mathcal{L}$ -module of level ℓ . Then there exists a unique $\mathcal{L}$ -homomorphism ψ from V to $\langle U \rangle$, sending 1_W to $\mathbf{1}$. Now, we have all the conditions assumed in Theorem 2.9. $\square$

Notice that vertex algebras are analogous to unital commutative associative algebras. It is a well known fact that any unital commutative associative algebra without zero divisor can be embedded into a field, namely a simple (and irreducible) commutative associative algebra. In view of this we formulate the following:

Conjecture 3.30. *Every nondegenerate vertex algebra can be embedded into a simple (irreducible) vertex algebra.*

4 Quantum vertex algebras associated with associative algebras of Zamolodchikov-Faddeev type

In this section we construct a family of quantum vertex algebras closely related to Zamolodchikov-Faddeev algebras.

Let H be a vector space equipped with a bilinear form $\langle \cdot, \cdot \rangle$ and let $\mathcal{S}(x) : H \otimes H \rightarrow H \otimes H \otimes \mathbb{C}[[x]]$ be a linear map. Recall from the end of Section 2 the notions of $(H, \mathcal{S})$ -module, vacuum-like vector and vacuum $(H, \mathcal{S})$ -module.

Set $\mathcal{S}(x) = \sum_{n \geq 0} S_n x^n$ where $S_n \in \text{End}(H \otimes H)$. Denote by $A(H, S_0)$ the quotient algebra of the (free) tensor algebra $T(H \otimes t^{-1}\mathbb{C}[t^{-1}])$ modulo the following relations:

$$(a \otimes t^{-m})(b \otimes t^{-n}) - \sum_{i=1}^r (b^{(r)} \otimes t^{-n})(a^{(r)} \otimes t^{-m}) = 0 \quad (4.1)$$

for $a, b \in H$, $m, n \geq 1$, where $S_0(b \otimes a) = \sum_{i=1}^r b^{(r)} \otimes a^{(r)}$. Notice that $A(H, 1)$ is the symmetric algebra over $H \otimes t^{-1}\mathbb{C}[t^{-1}]$ while $A(H, -1)$ is the exterior algebra.

Lemma 4.1. *Let H be a vector space equipped with a bilinear form $\langle \cdot, \cdot \rangle$ and let $\mathcal{S}(x)$ be a linear map from $H \otimes H$ to $H \otimes H \otimes \mathbb{C}[[x]]$. Let W be any vacuum $(H, \mathcal{S})$ -module with a vacuum-like vector e as its generator. For $k \geq 0$, let $W[k]$ be the span of the vectors*

$$a^{(1)}(-m_1) \cdots a^{(r)}(-m_r)e \quad (4.2)$$

for $r \geq 0$, $a^{(1)}, \dots, a^{(r)} \in H$, $m_i \geq 1$ with $m_1 + \cdots + m_r \leq k$. Then $W[k]$ for $k \geq 0$ form an increasing filtration of W such that for $a \in H$, $m, k \in \mathbb{Z}$,

$$a(m)W[k] \subset W[k - m], \quad (4.3)$$

$$a(m)W[k] \subset W[k - m - 1] \quad \text{whenever } m \geq 0, \quad (4.4)$$

where by convention we set $W[k] = 0$ for $k < 0$. Furthermore, $\text{gr}W = \coprod_{k \geq 0} W[k]/W[k-1]$ is an $A(H, S_0)$ -module with $e \in W[0] = W[0]/W[-1]$ as a generator, where

$$a(-m)(w + W[k-1]) = a(-m)w + W[k+m-1] \in W[k+m]/W[k+m-1]$$

for $a \in H$, $m \geq 1$, $w + W[k-1] \in W[k]/W[k-1]$ with $k \geq 0$.

Proof. From definition, we have $W[0] = \mathbb{C}e$ and by convention $W[k] = 0$ for $k < 0$. By definition we have

$$a(m)W[k] \subset W[k - m] \quad \text{for } a \in H, m < 0. \quad (4.5)$$

We next prove that

$$a(m)W[k] \subset W[k - m - 1] \quad \text{for } a \in H, m \geq 0. \quad (4.6)$$

For $a, b \in H$, as $\mathcal{S}(x)(b \otimes a) \in H \otimes H \otimes \mathbb{C}[[x]]$, we have

$$\iota_{x_2, x_1} \mathcal{S}(x_2 - x_1)(b \otimes a) = \sum_{r=1}^s \sum_{i \geq 0} \sum_{j=0}^i \beta_{rij} (b^{(r)} \otimes a^{(r)} \otimes x_2^{i-j} x_1^j),$$

where $\beta_{rij} \in \mathbb{C}$, $b^{(i)}, a^{(i)} \in H$. For $m, n \in \mathbb{Z}$, $w \in W$, from relation (2.34) we have

$$a(m)b(n)w = \sum_{r=1}^s \sum_{i \geq 0} \sum_{j=0}^i \beta_{rij} b^{(r)}(n + i - j) a^{(r)}(m + j) w + \langle a, b \rangle \delta_{m+n+1, 0} w. \quad (4.7)$$

As $W[0] = \mathbb{C}e$, from the assumption we have

$$a(m)W[0] = 0 = W[-m - 1] \quad \text{for } a \in H, m \geq 0.$$

Now, (4.6) follows immediately from induction (on k) and (4.7).

From (4.5) and (4.6), $\cup_{k \geq 0} W[k]$ is closed under the action of $a(m)$ for $a \in H$, $m \in \mathbb{Z}$. As $W = T(H \otimes \mathbb{C}[t, t^{-1}])e$, we have $W = \cup_{k \geq 0} W[k]$. Thus $W[k]$ for $k \geq 0$ form an increasing filtration of W . Using (4.7), (4.5) and (4.6) we have

$$\begin{aligned} & \left(a(-m)b(-n) - \sum_{r=1}^s \beta_{r00} b^{(r)}(-n) a^{(r)}(-m) \right) W[k] \\ &= \sum_{r=1}^s \sum_{i \geq 1} \sum_{j=0}^i \beta_{rij} b^{(r)}(-n + i - j) a^{(r)}(-m + j) W[k] \\ &\subset W[k + m + n - 1] \end{aligned}$$

for $m, n \geq 1$. Notice that

$$S_0(b \otimes a) = \sum_{r=1}^s \beta_{r00} (b^{(r)} \otimes a^{(r)}).$$

Now, the second assertion is clear. □

We also have:

Corollary 4.2. *Let $H, \langle \cdot, \cdot \rangle, \mathcal{S}$ and W, e be given as in Lemma 4.1. Then*

$$a^{(1)}(n_1) \cdots a^{(r)}(n_r)e = 0 \quad (4.8)$$

whenever $n_1 + \cdots + n_r \geq 0$ for $r \geq 1$, $a^{(i)} \in H$, $n_i \in \mathbb{Z}$.

Proof. If $n_1 + \cdots + n_r \geq 0$, we have $n_i \geq 0$ for some i . Using (4.3) and (4.4) we have

$$a^{(1)}(n_1) \cdots a^{(r)}(n_r)e \in W[-n_1 - \cdots - n_r - 1] = 0,$$

proving (4.8). □

Let $H, \langle \cdot, \cdot \rangle$, and $\mathcal{S}$ be given as in Lemma 4.1. Let $T(H \otimes \mathbb{C}[t, t^{-1}])^+$ be the subspace spanned by the vectors

$$(a^{(1)} \otimes t^{n_1}) \cdots (a^{(r)} \otimes t^{n_r})$$

for $r \geq 1$, $a^{(i)} \in H$, $n_i \in \mathbb{Z}$ with $n_1 + \cdots + n_r \geq 0$. Set

$$J = T(H \otimes \mathbb{C}[t, t^{-1}])T(H \otimes \mathbb{C}[t, t^{-1}])^+, \quad (4.9)$$

a left ideal of $T(H \otimes \mathbb{C}[t, t^{-1}])$. We then set

$$\tilde{V}(H, \mathcal{S}) = T(H \otimes \mathbb{C}[t, t^{-1}])/J, \quad (4.10)$$

a left $T(H \otimes \mathbb{C}[t, t^{-1}])$ -module. Clearly, $\tilde{V}(H, \mathcal{S})$ is cyclic on the vector $\tilde{1} = 1 + J$ where $a(n)\tilde{1} = 0$ for $a \in H$, $n \geq 0$. Furthermore, for $a \in H$, $w \in \tilde{V}(H, \mathcal{S})$, we have

$$a(m)w = 0 \quad \text{for } m \text{ sufficiently large.} \quad (4.11)$$

Now we define $V(H, \mathcal{S})$ to be the quotient $T(H \otimes \mathbb{C}[t, t^{-1}])$ -module of $\tilde{V}(H, \mathcal{S})$ modulo the following relations:

$$a(x_1)b(x_2)w - \sum_{i=1}^r \iota_{x_2, x_1}(f_i(x_2 - x_1))b^{(i)}(x_2)a^{(i)}(x_1)w = x_2^{-1}\delta\left(\frac{x_1}{x_2}\right)\langle a, b \rangle w \quad (4.12)$$

for $a, b \in H$, $w \in \tilde{V}(H, \mathcal{S})$, where $\mathcal{S}(x)(b \otimes a) = \sum_{i=1}^r b^{(i)} \otimes a^{(i)} \otimes f_i(x)$. Denote by $\mathbf{1}$ the image of $\tilde{1}$ in $V(H, \mathcal{S})$. From definition, $V(H, \mathcal{S})$ is a vacuum $(H, \mathcal{S})$ -module with vacuum-like vector $\mathbf{1}$ as its generator. In view of Corollary 4.2 we immediately have:

Proposition 4.3. *Let H be a vector space equipped with a bilinear form $\langle \cdot, \cdot \rangle$ and let $\mathcal{S}(x) : H \otimes H \rightarrow H \otimes H \otimes \mathbb{C}[[x]]$ be a linear map. The above constructed $(H, \mathcal{S})$ -module $V(H, \mathcal{S})$ is a universal vacuum $(H, \mathcal{S})$ -module.*

Combining Propositions 4.3 and 2.18 and Corollary 3.20 we immediately obtain:

Theorem 4.4. *Let H be a finite-dimensional vector space over $\mathbb{C}$ equipped with a bilinear form $\langle \cdot, \cdot \rangle$ and let $\mathcal{S}(x) : H \otimes H \rightarrow H \otimes H \otimes \mathbb{C}[[x]]$ be a linear map with $\mathcal{S}(0) = 1$. Assume that $V(H, \mathcal{S})$ is of PBW type in the sense that the $S(H \otimes t^{-1}\mathbb{C}[t^{-1}])$ -module $\text{gr}V(H, \mathcal{S}) = \coprod_{k \geq 0} V(H, \mathcal{S})[k]/V(H, \mathcal{S})[k-1]$, obtained in Lemma 4.1, is a free module. Then there exists a unique weak quantum vertex structure on $V(H, \mathcal{S})$ with $\mathbf{1}$ as the vacuum vector such that*

$$Y(a(-1)\mathbf{1}, x) = a(x) \quad \text{for } a \in H. \quad (4.13)$$

Furthermore, $V(H, \mathcal{S})$ is a nondegenerate quantum vertex algebra.

Notice that this theorem implies that if $V(H, \mathcal{S})$ is of PBW type, $\mathcal{S}(x)$ is necessarily a unitary rational quantum Yang-Baxter operator. We believe that if $\mathcal{S}(x)$ is a unitary rational quantum Yang-Baxter operator, $V(H, \mathcal{S})$ is of PBW type. Next we consider a special family, confirming our belief.

For the rest of this section we fix a finite-dimensional vector space U and an (invertible) element

$$R(x) = \sum_{n \geq 0} R_n x^n \in (\text{End}U)[[x]] \quad (4.14)$$

such that $R_i R_j = R_j R_i$ for $i, j \geq 0$ with $R_0 (= R(0))$ invertible. Then we have $R(x_1)R(x_2) = R(x_2)R(x_1)$.

Define $R^*(x) \in (\text{End}U^*)[[x]]$ by

$$\langle R^*(x)u^*, u \rangle = \langle u^*, R^{-1}(x)u \rangle \quad (4.15)$$

for $u \in U$, $u^* \in U^*$, where $\langle \cdot, \cdot \rangle$ is the standard pairing between U^* and U . Then

$$\langle R^*(x)u^*, R(x)u \rangle = \langle u^*, u \rangle \quad (4.16)$$

for $u \in U$, $u^* \in U^*$.

Set

$$H = U \oplus U^* \quad (4.17)$$

and equip H with the skew-symmetric bilinear form $\langle \cdot, \cdot \rangle$ defined by

$$\langle u + u^*, v + v^* \rangle = \langle u^*, v \rangle - \langle v^*, u \rangle \quad (4.18)$$

for $u, v \in U$, $u^*, v^* \in U^*$. Define $\mathcal{S}(x) : H \otimes H \rightarrow H \otimes H \otimes \mathbb{C}[[x]]$ by

$$\mathcal{S}(x)(u \otimes v) = R(-x)u \otimes R^{-1}(x)v, \quad (4.19)$$

$$\mathcal{S}(x)(u^* \otimes v^*) = (R^*)^{-1}(-x)u^* \otimes R^*(x)v^*, \quad (4.20)$$

$$\mathcal{S}(x)(v^* \otimes u) = R^*(-x)v^* \otimes R(x)u, \quad (4.21)$$

$$\mathcal{S}(x)(u \otimes v^*) = R^{-1}(-x)u \otimes (R^*)^{-1}(x)v^* \quad (4.22)$$

for $u, v \in U$, $u^*, v^* \in U^*$. We are going to show that if $R_0 = 1$, the universal vacuum $(H, \mathcal{S})$ -module $V(H, \mathcal{S})$ is of PBW type. Our strategy is to use certain vertex operator algebras associated to Heisenberg Lie algebras.

Remark 4.5. Consider the Heisenberg Lie algebra $\mathcal{H} = (U \oplus U^*) \otimes \mathbb{C}[t, t^{-1}] \oplus \mathbb{C}c$, where c is central and

$$\begin{aligned} [u \otimes t^m, v \otimes t^n] &= 0 = [u^* \otimes t^m, v^* \otimes t^n], \\ [u^* \otimes t^m, u \otimes t^n] &= \langle u^*, u \rangle \delta_{m+n+1, 0} c \end{aligned}$$

for $u, v \in U$, $u^*, v^* \in U^*$, $m, n \in \mathbb{Z}$. In terms of generating function $a(x) = \sum_{n \in \mathbb{Z}} (a \otimes t^n) x^{-n-1}$ for $a \in H$, we have

$$u(x_1)v(x_2) - v(x_2)u(x_1) = 0, \quad (4.23)$$

$$u^*(x_1)v^*(x_2) - v^*(x_2)u^*(x_1) = 0, \quad (4.24)$$

$$u^*(x_1)v(x_2) - v(x_2)u^*(x_1) = \langle u^*, v \rangle x_2^{-1} \delta\left(\frac{x_1}{x_2}\right) \quad (4.25)$$

for $u, v \in U$, $u^*, v^* \in U^*$. Set

$$\mathcal{H}^{\geq 0} = (U \oplus U^*) \otimes \mathbb{C}[t] \oplus \mathbb{C}c, \quad \mathcal{H}^- = (U \oplus U^*) \otimes t^{-1}\mathbb{C}[t^{-1}].$$

Consider $\mathbb{C}$ as an $\mathcal{H}^{\geq 0}$ -module with c acting as identity and with $(U \oplus U^*) \otimes \mathbb{C}[t]$ acting trivially. Form the induced $\mathcal{H}$ -module

$$V_{\mathcal{H}} = U(\mathcal{H}) \otimes_{U(\mathcal{H}^{\geq 0})} \mathbb{C}. \quad (4.26)$$

It is well known that $V_{\mathcal{H}}$ is an irreducible $\mathcal{H}$ -module and that $V_{\mathcal{H}}$ is naturally a simple conformal vertex algebra, which we shall use.

The vertex algebra $V_{\mathcal{H}}$ satisfies the following universal property:

Lemma 4.6. *Let K be any vertex algebra and let $\psi : U \oplus U^* \rightarrow K$ be any linear map such that*

$$\begin{aligned} [Y(\psi(u), x_1), Y(\psi(v), x_2)] &= 0 = [Y(\psi(u^*), x_1), Y(\psi(v^*), x_2)], \\ [Y(\psi(u^*), x_1), Y(\psi(u), x_2)] &= \langle u^*, u \rangle x_2^{-1} \delta\left(\frac{x_1}{x_2}\right) \end{aligned}$$

for $u, v \in U$, $u^*, v^* \in U^*$. Then ψ can be extended uniquely to a vertex algebra homomorphism from $V_{\mathcal{H}}$ to K .

Proof. From the assumption on ψ , K is an $\mathcal{H}$ -module with c acting as identity and with $a \otimes t^n$ acting as $\psi(a)_n$ for $a \in U \oplus U^*$, $n \in \mathbb{Z}$. We have $c \cdot \mathbf{1} = \mathbf{1}$ and $\mathcal{H}^{\geq 0} \mathbf{1} = 0$. Then there exists an $\mathcal{H}$ -module homomorphism f from $V_{\mathcal{H}}$ to K , sending the vacuum vector of $V_{\mathcal{H}}$ to the vacuum vector of K . Since $U \oplus U^*$ generates $V_{\mathcal{H}}$ as a vertex algebra, it follows that f is a vertex algebra homomorphism. $\square$

Recall from Remark 3.11 that Now, we have a vertex algebra $(\mathbb{C}[[t]], -d/dt)$ where

$$Y(f(t), x)g(t) = (e^{-x(d/dt)} f(t)) g(t) = f(t-x)g(t) \quad (4.27)$$

for $f(t), g(t) \in \mathbb{C}[[t]]$.

Lemma 4.7. *There exist unique vertex algebra homomorphisms $\Phi_R^\pm$ from $V_{\mathcal{H}}$ to the tensor product vertex algebra $\hat{V}_{\mathcal{H}} = V_{\mathcal{H}} \otimes \mathbb{C}[[t]]$ such that*

$$\Phi_R^\pm(u) = R^{\pm 1}(t)u, \quad \Phi_R^\pm(u^*) = (R^*)^{\pm 1}(t)u^* \quad (4.28)$$

for $u \in U$, $u^* \in U^*$.

Proof. Since U, U^* generate $V_{\mathcal{H}}$ as a vertex algebra, the uniqueness is clear. Because of the obvious symmetry we shall only prove the existence of Φ_R^+ . For $u, v \in U$, $u^*, v^* \in U^*$, we have

$$\begin{aligned} [\hat{Y}(R(t)u, x_1), \hat{Y}(R(t)v, x_2)] &= [Y(R(t-x_1)u, x_1), Y(R(t-x_2)v, x_2)] = 0, \\ [\hat{Y}(R^*(t)u^*, x_1), \hat{Y}(R^*(t)v^*, x_2)] &= [Y(R^*(t-x_1)u^*, x_1), Y(R^*(t-x_2)v^*, x_2)] = 0, \end{aligned}$$

and

$$\begin{aligned} &[\hat{Y}(R^*(t)u^*, x_1), \hat{Y}(R(t)u, x_2)] \\ &= [Y(R^*(t-x_1)u^*, x_1), Y(R(t-x_2)u, x_2)] \\ &= \langle R^*(t-x_1)u^*, R(t-x_2)u \rangle x_2^{-1} \delta\left(\frac{x_1}{x_2}\right) \\ &= \langle R^*(t-x_1)u^*, R(t-x_1)u \rangle x_2^{-1} \delta\left(\frac{x_1}{x_2}\right) \\ &= \langle u^*, u \rangle x_2^{-1} \delta\left(\frac{x_1}{x_2}\right). \end{aligned}$$

In view of Lemma 4.6, there exists a vertex algebra homomorphism Φ_R^+ from $V_{\mathcal{H}}$ to $\hat{V}_{\mathcal{H}}$ with the desired properties. $\square$

Lemma 4.8. *Let V be any vertex algebra. Equip $V \otimes \mathbb{C}[[t]]$ with the tensor product vertex algebra structure, where $\mathbb{C}[[t]]$ is viewed as a vertex algebra as above. A linear map $\Delta(t) : V \rightarrow V \otimes \mathbb{C}[[t]]$ is a vertex algebra homomorphism if and only if the following relations hold for $v \in V$:*

$$\Delta(t)\mathbf{1} = \mathbf{1} (= \mathbf{1} \otimes 1), \quad (4.29)$$

$$\Delta(t)Y(v, x) = Y(\Delta(t-x)v, x)\Delta(t). \quad (4.30)$$

Proof. Denote by $\hat{Y}$ the vertex operator map of the tensor product vertex algebra $V \otimes \mathbb{C}[[t]]$. Note that for $a \in V$, $f(t) \in \mathbb{C}[[t]]$, we have

$$\hat{Y}(a \otimes f(t), x) = Y(a, x)f(t-x).$$

For $u, v \in V$, we have

$$\hat{Y}(\Delta(t)u, x)\Delta(t)v = Y(\Delta(t-x)u, x)\Delta(t)v, \quad (4.31)$$

$$\Delta(t)\hat{Y}(u, x)v = \Delta(t)Y(u, x)v. \quad (4.32)$$

Then the assertion follows immediately. $\square$

Proposition 4.9. *Assume that $R_0 = R(0) = 1$. There exist unique elements*

$$\Delta_R^\pm(x) \in \text{Hom}_{\mathbb{C}}(V_{\mathcal{H}}, V_{\mathcal{H}} \otimes \mathbb{C}[[x]]) \subset (\text{End}_{\mathbb{C}} V_{\mathcal{H}})[[x]]$$

such that

$$\Delta_R^\pm(x)u = R^{\pm 1}(x)u, \quad \Delta_R^\pm(x)u^* = (R^*)^{\pm 1}(x)u^* \quad \text{for } u \in U, u^* \in U^*, \quad (4.33)$$

$$\Delta_R^\pm(x)\mathbf{1} = \mathbf{1}, \quad (4.34)$$

$$\Delta_R^\pm(x)Y(v, x_1) = Y(\Delta_R^\pm(x - x_1)v, x_1)\Delta_R^\pm(x) \quad \text{for } v \in V_{\mathcal{H}}, \quad (4.35)$$

$$\Delta_R^\pm(x_1)\Delta_R^\pm(x_2) = \Delta_R^\pm(x_2)\Delta_R^\pm(x_1), \quad \Delta_R^\pm(x_1)\Delta_R^\mp(x_2) = \Delta_R^\mp(x_2)\Delta_R^\pm(x_1), \quad (4.36)$$

$$\Delta_R^\pm(x)\Delta_R^\mp(x) = 1. \quad (4.37)$$

For $u \in U, u^* \in U^*$, set

$$u_R(x) = Y(u, x)\Delta_R^+(x) = u(x)\Delta_R^+(x), \quad (4.38)$$

$$u_R^*(x) = Y(u^*, x)\Delta_R^-(x) = u^*(x)\Delta_R^-(x). \quad (4.39)$$

Then $V_{\mathcal{H}}$ is a vacuum $(H, \mathcal{S})$ -module of PBW type.

Proof. In view of Lemma 4.8, such linear maps $\Delta_R^\pm(t)$ if exist are vertex algebra homomorphisms (from $V_{\mathcal{H}}$ to $V_{\mathcal{H}} \otimes \mathbb{C}[[t]]$). As $U \oplus U^*$ generates $V_{\mathcal{H}}$ as a vertex algebra, the uniqueness is clear. On the other hand, by Lemma 4.7, we have vertex algebra homomorphisms $\Phi_R^\pm$ from $V_{\mathcal{H}}$ to $V_{\mathcal{H}} \otimes \mathbb{C}[[t]]$ such that $\Phi_R^\pm(u) = R^{\pm 1}(t)u$ for $u \in U$ and $\Phi_R^\pm(u^*) = (R^*)^{\pm 1}(t)u^*$ for $u^* \in U^*$. Let us alternatively denote the map $\Phi_R^\pm$ by $\Delta_R^\pm(t)$. Then $\Delta_R^\pm(t)\mathbf{1} = \mathbf{1}$ and the relation (4.35) holds. For $u \in U$, we have

$$\Delta_R^\pm(x_1)\Delta_R^\pm(x_2)u = R^{\pm 1}(x_1)R^{\pm 1}(x_2)u = R^{\pm 1}(x_2)R^{\pm 1}(x_1)u = \Delta_R^\pm(x_2)\Delta_R^\pm(x_1)u.$$

From $R(x_1)R(x_2) = R(x_2)R(x_1)$ we have $R^*(x_1)R^*(x_2) = R^*(x_2)R^*(x_1)$, so that

$$\Delta_R^\pm(x_1)\Delta_R^\pm(x_2)u^* = (R^*)^{\pm 1}(x_1)R^*(x_2)u^* = R^*(x_2)R^*(x_1)u^* = \Delta_R^\pm(x_2)\Delta_R^\pm(x_1)u^*$$

for $u^* \in U^*$. Suppose

$$\Delta_R^\pm(x_1)\Delta_R^\pm(x_2)w = \Delta_R^\pm(x_2)\Delta_R^\pm(x_1)w$$

for some $w \in V_{\mathcal{H}}$. For any $a \in U \oplus U^*$, using (4.35) we have

$$\begin{aligned} \Delta_R^\pm(x_1)\Delta_R^\pm(x_2)Y(a, x)w &= Y(\Delta_R^\pm(x_1 - x)\Delta_R^\pm(x_2 - x)a, x)\Delta_R^\pm(x_1)\Delta_R^\pm(x_2)w \\ &= Y(\Delta_R^\pm(x_2 - x)\Delta_R^\pm(x_1 - x)a, x)\Delta_R^\pm(x_2)\Delta_R^\pm(x_1)w \\ &= \Delta_R^\pm(x_2)\Delta_R^\pm(x_1)Y(a, x)w. \end{aligned}$$

Since $U \oplus U^*$ generates $V_{\mathcal{H}}$ as a vertex algebra, it follows that

$$\Delta_R^\pm(x_1)\Delta_R^\pm(x_2)w = \Delta_R^\pm(x_2)\Delta_R^\pm(x_1)w \quad \text{for all } w \in V_{\mathcal{H}}.$$

This proves the first identity of (4.36). The second identity and (4.37) follow from the same argument.

For $u, v \in U$, using (4.35) and (4.36) we have

$$\begin{aligned}
& u_R(x_1)v_R(x_2) \\
&= Y(u, x_1)\Delta_R^+(x_1)Y(v, x_2)\Delta_R^+(x_2) \\
&= Y(u, x_1)Y(\Delta_R^+(x_1 - x_2)v, x_2)\Delta_R^+(x_1)\Delta_R^+(x_2) \\
&= Y(\Delta_R^+(x_1 - x_2)v, x_2)Y(u, x_1)\Delta_R^+(x_2)\Delta_R^+(x_1) \\
&= Y(R(x_1 - x_2)v, x_2)\Delta_R^+(x_2)Y(R^{-1}(x_2 - x_1)u, x_1)\Delta_R^+(x_1).
\end{aligned}$$

Similarly, for $u^*, v^* \in U^*$, we have

$$\begin{aligned}
& u_R^*(x_1)v_R^*(x_2) \\
&= Y(u^*, x_1)\Delta_R^-(x_1)Y(v^*, x_2)\Delta_R^-(x_2) \\
&= Y(u^*, x_1)Y(\Delta_R^-(x_1 - x_2)v^*, x_2)\Delta_R^-(x_1)\Delta_R^-(x_2) \\
&= Y(\Delta_R^-(x_1 - x_2)v^*, x_2)Y(u^*, x_1)\Delta_R^-(x_2)\Delta_R^-(x_1) \\
&= Y((R^*)^{-1}(x_1 - x_2)v^*, x_2)\Delta_R^-(x_2)Y(R^*(x_2 - x_1)u^*, x_1)\Delta_R^-(x_1).
\end{aligned}$$

We also have for $u \in U, u^* \in U^*$

$$\begin{aligned}
& u_R(x_1)v_R^*(x_2) - (R^*(x_1 - x_2)v^*)_R(x_2)(R(x_2 - x_1)u)_R(x_1) \\
&= Y(u, x_1)\Delta_R(x_1)Y(v^*, x_2)\Delta_R^-(x_2) \\
&\quad - Y(\Delta_R^+(x_1 - x_2)v^*, x_2)\Delta_R^-(x_2)Y(\Delta_R^+(x_2 - x_1)u, x_1)\Delta_R^+(x_1) \\
&= Y(u, x_1)Y(\Delta_R^+(x_1 - x_2)v^*, x_2)\Delta_R^+(x_1)\Delta_R^-(x_2) \\
&\quad - Y(\Delta_R^+(x_1 - x_2)v^*, x_2)Y(u, x_1)\Delta_R^+(x_1)\Delta_R^-(x_2) \\
&= Y(u, x_1)Y(R^*(x_1 - x_2)v^*, x_2)\Delta_R^+(x_1)\Delta_R^-(x_2) \\
&\quad - Y(R^*(x_1 - x_2)v^*, x_2)Y(u, x_1)\Delta_R^+(x_1)\Delta_R^-(x_2) \\
&= \langle u, R^*(x_1 - x_2)v^* \rangle x_2^{-1} \delta\left(\frac{x_1}{x_2}\right) \Delta_R^+(x_1)\Delta_R^-(x_2) \\
&= \langle u, R^*(0)v^* \rangle x_2^{-1} \delta\left(\frac{x_1}{x_2}\right) \Delta_R^+(x_2)\Delta_R^-(x_2) \\
&= \langle u, v^* \rangle x_2^{-1} \delta\left(\frac{x_1}{x_2}\right).
\end{aligned}$$

Substituting v^* with $(R^*)^{-1}(x_1 - x_2)v^*$ and u with $R^{-1}(x_2 - x_1)u$ and then exchanging

x_1 with x_2 we get

$$\begin{aligned}
& (R^{-1}(x_1 - x_2)u)_R(x_2)((R^*)^{-1}(x_2 - x_1)v^*)_R(x_1) - v_R^*(x_1)u_R(x_2) \\
&= \langle R^{-1}(x_1 - x_2)u, (R^*)^{-1}(x_2 - x_1)v^* \rangle x_1^{-1} \delta \left(\frac{x_2}{x_1} \right) \\
&= -\langle v^*, R(x_2 - x_1)R^{-1}(x_1 - x_2)u \rangle x_1^{-1} \delta \left(\frac{x_2}{x_1} \right) \\
&= -\langle v^*, u \rangle x_1^{-1} \delta \left(\frac{x_2}{x_1} \right). \tag{4.40}
\end{aligned}$$

This proves that $V_{\mathcal{H}}$ is an $(H, \mathcal{S})$ -module. Clearly, $\mathbf{1}$ is a vacuum-like vector.

On $V_{\mathcal{H}}$ we have an increasing filtration $\{E_k\}$ where for $k \geq 0$, E_k is spanned by the vectors

$$a^{(1)}(-m_1) \cdots a^{(r)}(-m_r) \mathbf{1}$$

for $r \geq 0$, $a^{(i)} \in H = U \oplus U^*$, $m_i \geq 1$ with $m_1 + \cdots + m_r \leq k$. For $a \in H$, $m, n \in \mathbb{Z}$, we have

$$\begin{aligned}
a(m)E_n &\subset E_{n+m}, \\
a(m)E_n &\subset E_{n-m-1} \quad \text{for } m \geq 0.
\end{aligned}$$

Furthermore, $\text{gr}_E(V_{\mathcal{H}})$ is a free $S(H \otimes t^{-1}\mathbb{C}[t^{-1}])$ -module.

For $a \in H$, set

$$a_R(x) = \sum_{n \in \mathbb{Z}} a_R(n) x^{-n-1}.$$

In the following we are going to prove that for each $k \geq 0$, E_k is also linearly spanned by the vectors

$$a_R^{(1)}(-m_1) \cdots a_R^{(r)}(-m_r) \mathbf{1}$$

for $r \geq 0$, $a^{(i)} \in H = U \oplus U^*$, $m_i \geq 1$ with $m_1 + \cdots + m_r \leq k$ and that

$$a_R(-m)w - a_R(-m)w \in E_{k+m-1} \tag{4.41}$$

for $a \in H$, $m \geq 1$, $w \in E_k$ with $k \geq 0$. Then it will follow immediately that $V_{\mathcal{H}}$ is a vacuum $(H, \mathcal{S})$ -module cyclic on the vacuum-like vector $\mathbf{1}$ and that $V_{\mathcal{H}}$ is of PBW type.

Since $E_0 = \mathbb{C}\mathbf{1}$ and $\Delta_R^{\pm}(x)\mathbf{1} = \mathbf{1}$, we have $\Delta_R^{\pm}(x)w - w = 0 \in E_{-1}$ for $w \in E_0$. For $a \in H = U \oplus U^*$ we have

$$\Delta_R^{\pm}(x)Y(a, x_1) = Y(\Delta_R^{\pm}(x - x_1)a, x_1)\Delta_R^{\pm}(x).$$

Set $\Delta_R^{\pm}(x)a = \sum_{r \geq 0} a^{(r)}x^r$ with $a^{(0)} = a$. For $m \in \mathbb{Z}$, we have

$$\Delta_R^{\pm}(x)a(m) = a(m)\Delta_R^{\pm}(x) + \sum_{r \geq 1} \sum_{i \geq 0} \binom{r}{i} (-1)^i a^{(r)}(m+i) x^{r-i} \Delta_R^{\pm}(x).$$

It follows from induction on k that for $w \in E_k$ with $k \geq 0$, we have

$$\Delta_R^\pm(x)w - w \in E_{k-1}[[x]] + xE_k[[x]]. \quad (4.42)$$

As $a_R(x) = a(x)\Delta_R^+(x)$ for $a \in U$ and $a_R(x) = a(x)\Delta_R^-(x)$ for $a \in U^*$, we always have

$$a_R(m)w - a(m)w \in \sum_{i \geq 0} a(m+i)E_{k-1} + \sum_{j \geq 1} a(m+j)E_k \subset E_{k-m-1}, \quad (4.43)$$

proving (4.41). From this we have

$$a_R^{(1)}(-m_1) \cdots a_R^{(r)}(-m_r)\mathbf{1} - a^{(1)}(-m_1) \cdots a^{(r)}(-m_r)\mathbf{1} \in E_{m_1+\cdots+m_r-1} \quad (4.44)$$

for $r \geq 0$, $a^{(i)} \in H$, $m_i \geq 1$. It follows from induction (on k) that E_k is linearly spanned by the vectors

$$a_R^{(1)}(-m_1) \cdots a_R^{(r)}(-m_r)\mathbf{1}$$

for $r \geq 0$, $a^{(i)} \in H = U \oplus U^*$, $m_i \geq 1$ with $m_1 + \cdots + m_r \leq k$. Therefore $V_{\mathcal{H}}$ as an $(H, \mathcal{S})$ -module is of PBW type. $\square$

Now we have:

Corollary 4.10. *Assume that $R_0 = R(0) = 1$. The universal vacuum $(H, \mathcal{S})$ -module $V(H, \mathcal{S})$ is of PBW type and $V(H, \mathcal{S})$ and $V_{\mathcal{H}}$ are isomorphic $(H, \mathcal{S})$ -modules.*

Proof. There exists a $T(H \otimes \mathbb{C}[t, t^{-1}])$ -module homomorphism f from $V(H, \mathcal{S})$ onto $V_{\mathcal{H}}$, sending the vacuum-like vector $\mathbf{1}$ to $\mathbf{1}$. Clearly, f reduces to an $S(H \otimes t^{-1}\mathbb{C}[t^{-1}])$ -module homomorphism $\bar{f}$ from $\text{gr}V(H, \mathcal{S})$ onto $\text{gr}V_{\mathcal{H}}$. Since $\text{gr}V_{\mathcal{H}}$ is a free $S(H \otimes t^{-1}\mathbb{C}[t^{-1}])$ -module and since both $\text{gr}V_{\mathcal{H}}$ and $\text{gr}V(H, \mathcal{S})$ are cyclic $S(H \otimes t^{-1}\mathbb{C}[t^{-1}])$ -modules, $\bar{f}$ is an isomorphism. Thus $\text{gr}V(H, \mathcal{S})$ is free. That is, $V(H, \mathcal{S})$ is of PBW type. From that $\bar{f}$ is an isomorphism, it follows that f is an isomorphism. $\square$

References

- [BK] B. Bakalov and V. Kac, Field algebras, *Internat. Math. Res. Notices* **3** (2003), 123-159.
- [B1] R. E. Borcherds, Vertex algebras, Kac-Moody algebras, and the Monster, *Proc. Natl. Acad. Sci. USA* **83** (1986), 3068-3071.
- [B2] R. E. Borcherds, Vertex algebras, in “*Topological Field Theory, Primitive Forms and Related Topics*” (Kyoto, 1996), edited by M. Kashiwara, A. Matsuo, K. Saito and I. Satake, Progress in Math., Vol. 160, Birkhäuser, Boston, 1998, 35-77.
- [B3] R. E. Borcherds, Quantum vertex algebras, preprint, arXiv:math.QA/9903038.

- [DL] C. Dong and J. Lepowsky, Generalized Vertex Algebras and Relative Vertex Operators, Progress in Math. **Vol. 112**, Birkhäuser, Boston, 1993.
- [DM] C. Dong and G. Mason, On quantum Galois theory, *Duke Math. J.* **86** (1997), 305-321.
- [D] V. G. Drinfeld, A new realization of Yangians and quantized affine algebras, *Soviet Math. Dokl.* **36** (1988), 212-216.
- [EK] P. Etingof and D. Kazhdan, Quantization of Lie bialgebras, V, *Selecta Math. (N.S.)* **6** (2000), 105-130.
- [F] L. D. Faddeev, Quantum completely integrable models in field theory, *Soviet Scientific Reviews Sect. C* **1** (1980), 107-155.
- [FKRW] E. Frenkel, V. Kac, A. Ratiu and W. Wang, $W_{1+\infty}$ and $W(gl_\infty)$ with central charge N , *Commun. Math. Phys.* **170** (1995), 337-357.
- [FR] E. Frenkel and N. Reshetikhin, Towards deformed chiral algebras, in: *Quantum Group Symposium, Proc. of 1996 Goslar conference*, ed. by H.-D. Doebner and V. K. Dobrev, Heron Press, Sofia, 1997, 27-42.
- [FHL] I. Frenkel, Y.-Z. Huang and J. Lepowsky, On axiomatic approaches to vertex operator algebras and modules, *Memoirs Amer. Math. Soc.* **104**, 1993.
- [FJ] I. Frenkel and N.-H. Jing, Vertex representations of quantum affine algebras, *Proc. Natl. Acad. Sci. USA* **85** (1988), 9373-9377.
- [FLM] I. Frenkel, J. Lepowsky and A. Meurman, *Vertex Operator Algebras and the Monster*, Pure and Appl. Math. **Vol. 134**, Academic Press, Boston, 1988.
- [FZ] I. B. Frenkel and Y. Zhu, Vertex operator algebras associated to representations of affine and Virasoro algebras, *Duke Math. J.* **66** (1992), 123-168.
- [K] V. G. Kac, *Vertex Algebras for Beginners*, University Lecture Series, Vol. 10, Amer. Math. Soc., 1997.
- [LL] J. Lepowsky and H.-S. Li, *Introduction to Vertex Operator Algebras and Their Representations*, Progress in Math. **227**, Birkhäuser, Boston, 2003.
- [Li1] H.-S. Li, Symmetric invariant bilinear forms on vertex operator algebras, *J. Pure Applied Algebra* **96** (1994), 279-297.
- [Li2] H.-S. Li, Local systems of vertex operators, vertex superalgebras and modules, *J. Pure Applied Algebra* **109** (1996), 143-195.
- [Li3] H.-S. Li, Axiomatic G_1 -vertex algebras, *Commun. Contemp. Math.* **5** (2003), 281-327.

- [Li4] H.-S. Li, Simple vertex operator algebras are nondegenerate, *J. Algebra* **267** (2003), 199-211.
- [Li5] H.-S. Li, Vertex algebras and vertex Poisson algebras, *Commun. Contemp. Math.* **6** (2004), 61-110.
- [Li6] H.-S. Li, G_1 -vertex algebras and their quasi modules, arXiv:math.QA/0502244.
- [MP] A. Meurman and M. Primc, Annihilating fields of standard modules of $\widetilde{\mathfrak{sl}(2, \mathbb{C})}$ and combinatorial identities, *Memoirs Amer. Math. Soc.* **652**, 1999.
- [MR] M. Mintchev and E. Ragoucy, Interplay between Zamolodchikov-Faddeev and reflection-transmission algebras, arXiv: math.QA/0306084.
- [Z] A. B. Zamolodchikov, Factorized S -matrices in two-dimensions as the exact solutions of certain relativistic quantum field theory models, *Ann. Phys.* **120** (1979), 253-291.